

AN ATLAS OF
REGIONAL DERMATOLOGY

This book is protected under the Berne Convention. It may not be reproduced by any means in whole or in part without permission. Application with regard to reproduction should be addressed to the Publishers.

An Atlas of Regional Dermatology

BY

G H PERCIVAL

M.D. Ph.D. F.R.C.P.E., D.P.H.

Grant Professor of Dermatology University of
Edinburgh Physician to the Skin Department,
Royal Infirmary Edinburgh

AND

T C DODDS

F.J.M.L.T. F.I.B.P. F.R.P.S.

Laboratory Supervisor Department of Pathology
University of Edinburgh Head of Photomicrography
Unit at Department of Pathology University of
Edinburgh

*Illustrated with 475 Clinical Subjects
in Full Colour*



E & S LIVINGSTONE LTD
EDINBURGH AND LONDON

1955

This book is protected under the Berne Convention. It may not be reproduced by any means in whole or in part without permission. Application with regard to reproduction should be addressed to the Publishers.

PREFACE

THIS Atlas is intended for use as an aid to the diagnosis of diseases of the skin. Its scope is limited to the presentation of representative examples of the common and less rare cutaneous disorders, as far as possible in their early developing and fully developed stages—and what is rare and what is common may vary with locality.

Many eruptions bear a close resemblance to one another and can only be differentiated by the recognition of minute details. In an Atlas the illustrations of such eruptions can be closely approximated thus enabling the observer to contrast and compare what is similar and in so doing note minor but important points of difference. Since this arrangement is based entirely on the morphology of the eruption, it cannot be conveniently employed in systematic textbooks on Dermatology.

Some skin diseases present a greater variety of eruptive elements than others, and consequently the clinical appearance of the eruption as a whole may differ to a considerable degree from case to case. Conversely diseases whose pathology and course are different may resemble each other closely as far as their skin manifestations are concerned. Such conditions naturally require a larger number of illustrations than do those which are more constant or distinctive in their appearances, and this applies to the rarer diseases such as Dermatitis Herpetiformis and Pemphigus, as well as to the more common ones such as the cutaneous cancers.

A regional arrangement of the Atlas has been chosen for several reasons, even if it involves some duplication. According to the site affected a given eruption may show variations from what is described as its normal appearance. Most skin diseases may occur on any part of the body surface, but the majority tend to be selective in the regions which they affect, even although such selectivity is seldom exclusive or uniform. In the clinic most diseases of the skin present themselves as regional problems.

In conclusion the Authors have pleasure in recording their indebtedness to the University of Edinburgh for the use of the transparencies reproduced in this Atlas. They also wish to acknowledge the help given by Dr D. A. Duthie in connection with the clinical photography and Dr Allen Lyell in the preparation of the index.

G. H. PERCIVAL,
T. C. DODDS.

CONTENTS

	AGE
HEAD AND NECK	1
HANDS AND ARMS	82
FEET AND LEGS	115
TRUNK AND GENERALISED ERUPTIONS	157
INDEX	259



FIG. 1

Impetigo Contagiosa

The lesions are superficial subcorneal bullae the contents of which have coagulated to form crusts. On the centre of the cheek crust has been removed to expose the raw base of bulla and the remains of the bulla roof are visible as scale bordering the lesion.



FIG. 2

Impetigo Contagiosa

The lesions are numerous and the crusts have coalesced.



FIG. 5

Impetigo Contagiosa

The lesions are scanty and discrete, and at first sight might suggest *acne vulgaris*. Their sudden onset, bullous character, and short duration serve as distinguishing features.



FIG. 6

Papulovesicular Erythema

The lesions may bear superficial resemblance to impetigo, acne, and rosacea, but they consist of *indurated papules*, and their duration is limited to a few weeks. The details of individual lesions are shown in Fig. 395.



FIG. 3

Impetigo Contagiosa

The same case as in Fig. 2. The crusts have been removed revealing the oozing bases of numerous bullae.



FIG. 4

Impetigo Contagiosa

The eruption consists of the glistening bases of numerous bullae and crusting is not a prominent feature. The lesions tend to be circinate in places, possibly suggesting *tinea circinata*. A more purely bullous character is the distinguishing feature and further confirmation could be obtained by microscopic examination of epidermal tags at the edge of the lesions.



FIG. 9

Herpes Simplex

The eruption is more extensive than usual the vesicles have become converted into crusts there is some resemblance to impetigo, but the grouping of the lesions is apparent.



FIG. 7

Herpes Simplex

The lesions consist of firm vesicles grouped in bunches. Acne lesions are present on the cheeks and chin



FIG. 8

Herpes Simplex

Herpes simplex of lip and cheek. The grouping of the firm vesicles and the limited duration of the eruption, usually five to ten days, differentiate it from impetigo. Recurrences are common, and the eruption may be bilateral.




FIG. 12

Herpes Zoster

Eruption on the sixth day. It is now crusted and slightly hemorrhagic. The upper eyelid has become affected and there is considerable edema of the corresponding cheek.



FIG. 13

Herpes Zoster

Eruption on the eighth day. The crusts have been removed revealing the raw upper surface of the dermis. Healing will be completed by scar formation.



FIG. 10

Herpes Zoster

The eruption has been present for three days. It consists of firm vesicles arranged in bunches on an area of erythema. The linear and unilateral distribution is apparent.



FIG. 11

Herpes Zoster

Eruption on the fifth day. It is unilateral, the vesicles, which are becoming purulent, are grouped as in herpes simplex but are more deeply situated in the skin.



FIG. 16A
Thrush Barber (Karlson type)



FIG. 16
Thrush Barber of Neck

In Figs. 16A, 16B indurated inflammatory nodules are prominent and distinguish the condition clinically from staphylococcal pyoderma, even when secondary staphylococcal folliculitis has developed. The fungus infection is slowly eradicated, through the development of tissue immunity. The condition is then self-healing and of limited duration. It lasts from six weeks to three months and may leave scars. In some cases its clinical appearance might suggest syphilis.



FIG. 14

Sycosis Barber (Staphylococcal)

As the case is an early one, pustulation is clearly seen. Later as a result of recurrent pustulation, the affected areas become persistently erythematous and slightly swollen, and in these circumstances pustules are less clearly seen.



FIG. 15

Sycosis Barber (Staphylococcal)

Pustulation is less in evidence. Chronic congestion and slight swelling of the skin are present together with a chronic eczema reaction. The case is of long standing.



FIG. 19

Acne Vulgaris

Posterior lesions are mingled with infected sebaceous cysts.



FIG. 20

Acne Vulgaris



FIG. 17

Acne Vulgaris

The lesions are composed of blackheads (comedones) and the papules and pustules which develop at the mouths of the follicles which are plugged by them.



FIG. 18

Acne Vulgaris

Pustules and elongated indolent superficial abscesses are present.



FIG. 23
Rosacea

This eruption has been present for a few years and bears some resemblance to lupus erythematosus, but the eruptive elements are papules and pustules which vary in amount from week to week.



FIG. 24
Rhinophyma

This condition may complicate rosacea or arise spontaneously. It is of slow development but permanent. It consists of hyperplasia of the sebaceous glands of the area, accompanied by an increase in the vascularity of the area. These changes manifest themselves in the form of soft tumour-like enlargement of the nose. Periodic pustulation may be a troublesome feature.



FIG 21

Roseacea

The lesions consist of persistent erythema and recurrent papules which may become purulent. The papules are not preceded by blackheads, and the disease occurs rarely if ever during adolescence.



FIG 22

Roseacea

The eruptive elements come and go on the blush area, and the appearance varies from week to week. This, together with the fact that the lesions are papules and pustules, differentiates it from lupus erythematosus.



FIG. 26

Papulosecretory Tuberculosis

The lesions consist of firm papules of varying size which persist for a number of weeks and eventually become necrotic. Scar formation may accompany healing. As a result of successive eruptions of papules the condition may persist for number of months. As is seen in subsequent illustrations, the amount of the eruption may vary considerably.



FIG. 27

Papulosecretory Tuberculosis

A more florid example than that shown in Fig. 26.



FIG. 25

Nodulocystic Syphilis

The nodular character of the eruption distinguishes it from rosacea, as does its comparatively short duration. Serological tests for syphilis are positive. The nose, particularly the alae nasi, is a not uncommon site for late secondary and early tertiary syphilitic lesions.



FIG. 30

Annular Sarcoid

This is an unusual configuration of a superficial sarcoid. The lesions consist of persistent superficial nodules arranged in annular form. The histopathology is typical. Any epidermal change is secondary to the infiltrate in the dermis, and should its presence in the form of scaling suggest *tinea circinata*, the absence of foreign elements in the scales will exclude the latter condition.



FIG. 31

Reticular Cell Sarcoma

The form of the lesion is an infiltrated plaque of gray outline and the diagnosis is only possible from the histopathology



FIG. 28

Papulonecrotic Tuberculide

The histology of the lesions is typical in this disease.



FIG. 29

Sarcoid

The lesion is an indolent, diffuse painless swelling which develops slowly and often has a violaceous colour. The diagnosis rests on the histopathology which is typical. Clinically a reticulum cell sarcoma might be considered.



FIG. 34

Adams-Schicron

Case of tuberous sclerosis the associated periorbital fibromas are shown in Fig. 227



FIG. 35

Adams-Schicron



FIG. 32

Adenoma Sebaceum

The lesions consist of small red nodules composed of hyperplastic sebaceous glands and vessels. It is usually associated with some degree of mental deficiency and may accompany tuberous sclerosis.



FIG. 33

Adenoma Sebaceum



FIG. 34

Adenoma Sebaceum

Case of tubercularosis the associated periorbital fibromata are shown in Fig. 227



FIG. 35

Adenoma Sebaceum



FIG. 32

Adenoma Sebaceum

The lesions consist of small red nodules composed of hyperplastic sebaceous glands and vessels. It is usually associated with some degree of mental deficiency and may accompany tuberous sclerosis.



FIG. 33

Adenoma Sebaceum



FIG. 38

Lupus Erythematosus

There are erythematous plaques on which telangiectatic vessels are visible. The absence of recurring papules and pustules, and the stationary character of the eruption, differentiate it from rosacea.



FIG. 39

Lupus Erythematosus

There are chronic discoid plaques with thick adherent masses of keratin on the surface. Atrophic scarring is also in evidence.



FIG 36

Lupus Erythematosus

The eruption is of recent origin and consists of erythematous plaques which are slightly raised. As in all varieties of lupus erythematosus, apple jelly nodules cannot be demonstrated by vitropression.



FIG 37

Lupus Erythematosus

The eruption consist of infiltrated discoid lesions which have a rough surface due to hyperkeratosis and plugging of the follicle mouths by keratin.



FIG. 38

Lupus Erythematosus

There are erythematous plaques on which telangiectatic vessels are visible. The absence of recurring papules and pustules, and the stationary character of the eruption, differentiate it from rosacea.



FIG. 39

Lupus Erythematosus

There are chronic discoid plaques with thick adherent masses of keratin on the surface. Atrophic scarring is also in evidence.



FIG 40

Lupus Vulgaris

This eruption is not complicated by secondary features such as scaling, crusting or ulceration. The eruption consists of a slightly swollen erythema which has a brownish tinge. Vitropression reveals "apple-jelly nodules" which differentiate it from lupus erythematosus. The absence of papules and pustules and week-to-week variation in appearance distinguish it from rosacea, and its long duration and the results of serological investigation from syphilis.



FIG 41

Lupus Vulgaris

A lesion of recent origin.



FIG. 42

Lupus Vulgaris

The clearly defined irregular margin of the plaques is evident and the brownish-red colour of the eruption is typical.



FIG. 43

Lupus Vulgaris

The eruption is composed of an aggregation of small tubercles situated in the dermis. There is some secondary scaling.



FIG. 44

Lupus Vulgaris

The disease has lasted for many years and has caused mutilation of the cartilages. The scarring, which is mingled with still active disease is due to treatment.



FIG. 45

Lupus Vulgaris



FIG. 46

Flexural Infective Eczema

Early flexural infective eczema with secondary pustulation and impetigo on surrounding skin.



FIG. 47

Flexural Infective Eczema

Early stage, showing fissuring at the depth of the fold.



FIG. 44

Lupus Vulgaris

The disease has lasted for many years and has caused mutilation of the cartilages. The scarring, which is mingled with still active disease, is due to treatment.



FIG. 45

Lupus Vulgaris



FIG. 50

Flexural Infective Eczema

Starting behind both ears as in Fig. 46, the reaction has extended to involve the whole scalp and has spread to the adjacent skin of the forehead and eyebrows.



FIG. 51

Flexural Infective Eczema

A retro-auricular flexural infective eczema which has spread to involve the scalp.



FIG. 48

Flexural Infective Eczema

To show the symmetry of the lesion on each side of the retro-auricular fold (see subsequent illustrations of other folds).



FIG. 49

Flexural Infective Eczema

The eczema reaction has spread from the retro-auricular fold to the adjacent scalp while the outer aspect of the pinna remains free



FIG. 54

Flexural Infective Eczema

Symmetrical involvement of the fold of the upper eyelid. Note the absence of edema and compare with contact eczema on the same area.



FIG. 55

Flexural Infective Eczema

The upper and lower lids and the canthi are involved. Note the absence of edema. The retro-auricular folds were affected first.



FIG. 52

Flexural Infective Eczema

Similar case to Fig. 51. Such cases are also included in the ill-defined and indefinite group of "Seborrhæic Eczema."




FIG. 53

Flexural Infective Eczema

This case illustrates one mode of origin of flexural infective eczema. There is a primary impetigo of the face, neck, and retro-auricular region. The impetigo later cleared under treatment leaving in its wake a persistent retro-auricular infective eczema.



FIG. 58

Contact Eczema

The reaction has been caused by an ingredient in lipstick.




FIG. 59

Contact Eczema

Positive patch test (lipstick)



FIG. 56

Contact Eczema

The reaction is due to sensitivity to an eye lotion. Note the edema associated with the reaction and compare with Fig. 55



FIG. 57

Contact Eczema

The reaction is due to sensitivity to atropin. Note the widespread edema.



FIG. 62
Infantile Eczema



FIG. 63
Infantile Eczema
The eruption has developed secondary impetigo



FIG. 60
Solar Eczema

Note the distribution of the reaction on the prominences of the face and the absence of oedema of eyelids. The condition might in some cases suggest rosacea or lupus erythematosus. Apart from the recognition that the eruption is an eczema reaction, the history of seasonal occurrence is a help in diagnosis.



FIG. 61
Solar Eczema

Note the involvement of the lobe of ear. The distribution of the eruption in this area is a help in diagnosis.



FIG. 65

Acne Keloid

This shows the early chronic recurrent pustular stage. The condition is staphylococcal folliculitis. An anatomical abnormality in a number of the hair follicles on the affected area, consisting of the incorporation of several hairs in single follicle, is predisposing factor



FIG. 66

Acne Keloid

Stage of hypertrophic scar formation. Due to the recurrent folliculitis results in the development of areas of granulation tissue which destroys the hairs on the area and which clinically resembles keloid.



FIG 64

Pityriasis Rubra Pilaris

The eruption is purely erythematous, and scaling is seldom a marked feature. The presence of the eruption on other areas may be necessary for a diagnosis. As a rule the histopathology is inconclusive as a diagnostic factor



FIG. 65
Acne keloid

This shows the early chronic recurrent pustular stage. The condition is staphylococcal folliculitis. A anatomical abnormality in a number of the hair follicles on the affected area, consisting of the incorporation of several hairs in single follicle, is a predisposing factor



FIG. 66
Acne keloid

Stage of hypertrophic scar formation. In due course the recurrent folliculitis results in the development of areas of granulation tissue which destroys the hairs on the area and which clinically resembles keloid.



FIG. 64

Pityriasis Rubra Pilaris

The eruption is purely erythematous, and scaling is seldom a marked feature. The presence of the eruption on other areas may be necessary for a diagnosis. As a rule the histopathology is inconclusive as a diagnostic factor.



FIG. 69

Lichen Simplex

This is sequel to patch of post-traumatic infective eczema. The skin is thickened and of leathery texture. The term neurodermatitis is also applied to such localized thickened patches.



FIG. 70

Thin Carpath



FIG. 67

Acne keloid

Stage of atrophy. The keloid-like mass eventually resolves itself and gives place to an atrophic scar.



FIG. 68

Xanthoma

Eruption of xanthomata in a case of *xanthoma tuberosum*. The lesions appear as firm yellowish nodules of varying size. These persist for a variable period and then disappear. They tend to erupt in successive crops, so that the eruption as a whole may be of long duration.



FIG. 73

Hyperkeratotic Nevus

The structure of this birthmark is hypertrophy of the stratum corneum. Lesions of this type are almost always linear and may cover large areas.



FIG. 74

Mixed Hyperkeratotic Nevus and Nevus Sebaceus

There is congenital hypertrophy of the stratum corneum and the sebaceous glands. A linear pattern is usual.



FIG. 71

Fixed Drug Eruption

A recurrent erythema which in an individual case always occurs on the same site. In this patient it is caused by phenolphthalein.




FIG. 72

Erysipoid

Chronic recurring erythematous eruption (erysipoid). The condition is affecting the cheek and upper lip. It resembles erysipela but toxic symptoms are absent. The condition is recurrent and the affected areas become somewhat thickened, presumably result of blockage in the cutaneous lymphatics.



FIG. 77

Nervus Suboculcus

In this case the lesions are single and do not show the usual linear pattern. The diagnosis can be made only by histological examination.



FIG. 78

Nervus Suboculcus



FIG. 75

Nævus Syringadenomatosus Papilliferus

In this tumour the hypertrophy is limited to the sweat glands and their ducts. The surface is frequently eroded in places, giving the impression of granulation tissue.




FIG. 76

Nævus Sebaceus

Nævi of the type shown in Figs. 75 and 76 can only be differentiated histologically



FIG. 77

Nemes Schöcs

In this case the lesions are single and do not show the usual linear pattern. The diagnosis can be made only by histological examination.



FIG. 78

Nemes Sebastes



FIG. 79

Nævus Syringadenomatosus Papilliferus

This nævus may present an eroded surface. Histological examination is necessary to distinguish accurately between it and the preceding nævus.



FIG. 80

Diffuse Alopecia (Post-febrile type)

In this case the fall of hair took place about six weeks after recovery from pneumonia. Regrowth is the rule.



FIG. 81

Diffuse Alopecia (Syphilitic type)

The occipital region usually best shows the incomplete alopecia of secondary syphilis.



FIG. 82

Circumscribed Alopecia (Alopecia Areata)

The clean shiny surface of the bald area is characteristic of the disease and differentiates it from tinea capitis.



FIG. 83

Circumscribed Alopecia (Pseudopelade of Brocq)

Note the irregular pattern this form of alopecia is permanent, and it develops and spreads slowly



FIG. 84

Circumscribed Alopecia (Folliculitis Decumbens)

The scanty pustulation is in evidence. It slowly produces a cicatrix with permanent loss of hair



FIG. 85

Circumscribed Alopecia (Folliculitis Decumbens)

The pustulation is profuse, resembling sycosis.



FIG. 82

Circumscribed Alopecia (Alopecia Areata)

The clean shiny surface of the bald area is characteristic of the disease and differentiates it from thin capitis.



FIG. 83

Circumscribed Alopecia (Pseudopelade of Brocq)

Note the irregular pattern—this form of alopecia is permanent, and it develops and spreads slowly




FIG. 84

Circumscribed Alopecia (Folliculitis Decalvans)

The scanty pustulation is in evidence. It slowly produces alopecia with permanent loss of hair



FIG. 85

Circumscribed Alopecia (Folliculitis Decalvans)

The pustulation is profuse, resembling sycomata.



FIG 86

Circumscribed Alopecia (Lupus Erythematosus)

Active erythematous areas are to be seen and these and the atrophic scars which result from them cause permanent destruction of the follicles.




FIG 87

Circumscribed Alopecia (Syphilitic)

A gummatous syphilide which has produced extensive cicatrization and permanent loss of hair



FIG. 88

Thinea Capitis

The partially bald patch is covered with thick matting of scales, and short broken and twisted hairs are present on the surface. This contrasts with the smooth clean bald patches of alopecia areata. Examination of the patient under Wood's light and microscopic examination of hairs should be carried out



FIG. 89

Thinea Kerion

The reaction to the fungus invasion is more definite in this case and pustulation is present. This is commencing kerion which became fully developed in the succeeding week.



FIG. 90

Thien Kerion

This shows a small well-developed kerion of the scalp. The appearance suggests that of an aggregation of small cutaneous abscesses.



FIG. 91

Favus Capitis

The hairs are matted with dry squameous crusts, an appearance liable to be mistaken for impetigo or eczema.



FIG. 92

Favus Capitis

The scalp is covered with friable squame crusts, and completely bald atrophic areas are to be seen.



FIG. 93

Favus Capitis

The greater part of the scalp has become atrophic and permanently denuded of hair. Masses of adherent friable squame crusts are present and few isolated cup-shaped scutula are to be seen.




FIG. 94

Favus Capitis

The crusting is composed of fungus elements and desiccated exudate. Permanent cicatricial baldness is present.



FIG. 95

Favus Capitis

As in the preceding examples scutula, massive crusting, scarring, and alopecia are present



FIG. 96

Village

The skin texture is normal and the change consists in the loss of the capacity to form pigment in certain areas. The adjacent areas are often hyperpigmented.



FIG. 97

Senile Evertsels

The skin is yellowish in colour hypertrophied, and the natural folds exaggerated. This is a normal occurrence on exposed areas, and is due to chemical changes in the collagen bundles of the dermis.



FIG. 98

Scleroderma

In addition to being sclerosed the affected area is depigmented. In some cases, particularly in the later stages when atrophy may have succeeded sclerosis, the appearance may suggest lichen sclerosus et atrophicus, and a histological examination may be necessary to establish a diagnosis. The hard texture of scleroderma serves to differentiate it from vitiligo. In calcinosis cutis the lesions are stony hard.



FIG. 99

Pelliodermis (Atrophicus Vasculare)

This is a slowly progressive atrophy associated with the development of telangiectasia, irregular hyperpigmentation and the whitish area of atrophic skin. It is not preceded by any inflammatory eruption. It may resemble the changes seen in long-standing lupus erythematosus, and a somewhat similar change may be present in dermatomyositis.



FIG. 100
Accidental Tattooing



FIG. 98
Scleroderma

In addition to being sclerosed the affected area is depigmented. In some cases, particularly in the later stages when atrophy may have succeeded sclerosis, the appearance may suggest lichen sclerosis et atrophicus, and a histological examination may be necessary to establish a diagnosis. The hard texture of scleroderma serves to differentiate it from vitiligo. In calcinosis cuts the lesions are stony hard.



FIG. 99
Pothodermia (Atrophia Vasculare)

This is a slowly progressive atrophy associated with the development of telangiectasia, irregular hyperpigmentation and thin whitish areas of atrophic skin. It is not preceded by any inflammatory eruption. It may resemble the changes seen in long-standing lupus erythematosus, and a somewhat similar change may be present in dermatomyositis.



FIG. 103
Capillary Angioma
Spider naevus type



FIG. 104
Capillary Angioma
Port-wine stain variety



FIG. 101
Fibroma



FIG. 102
Mixed Cell Nevus

The lesion consists of nevus cells, hair follicles, and sweat glands and ducts. This type of tumour may appear for the first time, or enlarge, about the menopause



FIG. 103
Capillary Angioma
Spider nevus type.



FIG. 104
Capillary Angioma
Port-wine stain variety



FIG 105
Capillary Angioma
"Port-wine stain" variety



FIG 106
Cavernous Angioma

The lesions of the forehead are superficial, that on the lid is superimposed on a deeper seated cavernous angioma.



FIG. 107
Verruca Etilica



FIG. 108
Xanthelasma



FIG 109
Atrocliform Nerve and Sebaceous Cyst



FIG. 110

Millers

The lesions are keratin-containing epidermoid cysts which can be stalked out with ease. They are of common occurrence in anidic ge. See colloid pseudomorphs.



FIG. 111

Millers

Many are infected and becoming pustular



FIG. 116

Cutaneous Horn and Squamous Cell Carcinoma

This lesion is an extension of the preceding one: the hypertrophied epidermis at the base of the horn having developed malignant tendencies.



FIG. 117

Squamous Cell Carcinoma

Microscopic examination is necessary for the correct diagnosis of cutaneous carcinomata. In early cases the clinical features are not sufficiently well defined to differentiate between the two types. The succeeding examples of the varying appearances which the two types may present illustrate this point.



FIG. 118

Basal Cell Carcinoma



FIG. 119

Squamous Cell Carcinoma

This has developed on an X ray scar produced in the treatment of a patch of lupus vulgaris. Active lupus is still present on the lobe of the ear.



FIG 116

Cutaneous Horn and Squamous Cell Carcinoma

This lesion is an extension of the preceding one the hypertrophied epidermis at the base of the horn having developed malignant tendencies.



FIG 117

Squamous Cell Carcinoma

Microscopic examination is necessary for the correct diagnosis of cutaneous carcinomata. In early cases the clinical features are not sufficiently well defined to differentiate between the two types. The succeeding examples of the varying appearances which the two types may present illustrate this point.



FIG. 122

Squamous Cell Carcinoma

This has developed on patch of *lepus vulgaris*.



FIG. 120

Squamous Cell Carcinoma

This has developed on an extensive area of lupus,
previously treated by X-rays.



FIG. 121

Basal Cell Carcinoma

Cylindroma type.



FIG. 14
Basal Cell Carcinoma
Cylindroma type.



FIG. 125
Multiple Basal Cell Carcinomas



FIG. 123

Squamous Cell Carcinoma

The same patient after treatment with X-rays.



FIG. 128
Basal Cell Carcinoma



FIG. 129
Squamous Cell Carcinoma



FIG. 126
Squamous Cell Carcinoma



FIG. 127
Cystic Basal Cell Carcinoma



FIG. 131

Xeroderma Pigmentosum

This shows the atrophy which is partly associated with the skin condition and partly due to the treatment of numerous carcinomata. The patient is aged twenty-four and the condition was first noticed at the age of four years.



FIG 130
Basal Cell Carcinoma



FIG. 134
Lentiginis Culi



FIG. 135
Myxoid Erythema

The eruption bears some clinical resemblance to nodulocutaneous myxoid, and the diagnosis is determined by the histopathology and serology




FIG 132

Sarcoma

The tumours are producing the appearance of a partial "turban tumour". Multiple basal cell carcinomata of the cylindroma type may show a similar clinical appearance.



FIG 133

Sarcoma

The eyelids and scalp are affected.



FIG. 138

Melanoma

Melanoma with confluent area in the centre.



FIG. 139

Malignant Melanoma



FIG 136

Mycosis Fungoides

In this example the infiltrate of reticulum cells does not extend deeply into the dermis, so that the lesion is not markedly tumour-like. The surface is eroded, and at first glance the condition bears some resemblance to an eczema reaction.



FIG 137

Malignant Melanoma



FIG. 142

Syphilitic Chancere

The lower lip is affected. Note the marked swelling of the tissues associated with the ulcer.



FIG. 143

Syphilitic Chancere

Lower lip. There is contrasting papuloqueous eruption on the chin.



FIG. 140

Blue Nevus

This is a benign melanoma in which the pigment-forming and pigment-containing cells are situated deep in the dermis




FIG. 141

Halo Nevus

A small benign melanoma with depigmentation in the surrounding skin.



FIG. 142

Syphilitic Chancere

The lower lip is affected. Note the marked swelling of the tissues associated with the ulcer



FIG. 143

Syphilitic Chancere

Lower lip. There is constricting papulosquamous eruption on the chin.



FIG 144

Lichen Planus

The lesions consist of papules which are identical with those which occur on the skin. The white appearance on the mucous membrane is due to the formation of a stratum granulosum and corneum, structures which are not normally present in the epithelium of the mucous membranes.



FIG 145

Lichen Planus

The network pattern is commonly seen. Lichen planus may affect the mucous membrane alone the skin remaining free from lesions.



FIG. 146
Lichen Planus

Isolated white papules are visible on the mucous membrane of the cheek.




FIG. 147
Lupus Erythematosus

The eruption consists of redness and adherent scaling of the lips. It may occur on the lips alone, unaccompanied by cutaneous lesions, and in these circumstances the



FIG 148

Leukoplakia

Small retrocommunal patches of leukoplakia are common. As in lichen planus, the white appearance is due to hyperkeratosis developing on the mucous membrane epithelium. In leukoplakia the change is not limited to the surface of papules but occurs as a diffuse plaque.



FIG 149

Leukoplakia

The hyperkeratosis is marked and the area is already showing malignant change.



FIG. 150
Leukoplakia



FIG. 151
Leukoplakia



FIG. 152
Contact Eczema

Note the large single area affected, the presence of formed vesicles, and the gradual fading of the reaction at the periphery without any defined edge.



FIG. 153
Contact Eczema

Note the large area involved by the eruption, the lack of any demarcation at its edge, and its vesicular character. The thick stratum corneum of the palm holds the initial crop of vesicles intact for a longer period than would be the case on other parts of the body. On the fingers the corneum forming the 'roof' of numerous vesicles has been removed, revealing the upper surface of the rete mucosum and, incidentally, causing the eruption to become much edematous in appearance.



FIG. 154

Post-traumatic Infective Eczema

Note the clearly defined edge of the individual lesions in the shape of circles or segments of circle and the multiplicity of lesions in close approximation. Exudation is apparent and is proceeding from vesicles which are now forming in an epidermis denuded of its stratum corneum as result of previous crops of vesicles. Compare the contour of the individual lesions with those of contact eczema.



FIG. 155

Post-traumatic Infective Eczema

Note the clearly defined margin of the eruption and the close proximity of adjacent patches.



FIG. 15

Contact Eczema

Note the large single area affected, the presence of formed vesicles, and the gradual fading of the reaction at the periphery without any defined edge



FIG. 153

Contact Eczema

Note the large area involved by the eruption, the lack of any demarcation at its edge and its vesicular character. The thick stratum corneum of the palm holds the initial crop of vesicles intact for a longer period than would be the case on other parts of the body. On the fingers the corneum forming the roofs of numerous vesicles has been removed revealing the upper surface of the rete mucosum and, incidentally, causing the eruption to become much redder in appearance.



FIG. 158

Eczema

This shows the effect of crude gas tar on the eczema reaction. The tar has been easily rubbed off the areas previously affected by the eczema reaction and they are clearly outlined by the medication. This indicates that although the patches appear to be healed, the cornium is not yet completely restored to normal.



FIG. 159

Flexural Infective Eczema

A flexural infective eczema of the elbow fold with chemical eczema on the upper arm due to the application



FIG. 156

Post-traumatic Infective Eczema

Note the sharply-defined edge of lesions and compare with contact eczema.



FIG. 157

Post-traumatic Infective Eczema

The eruption is in a scaly phase. It bears some superficial resemblance to psoriasis, but in the latter the scales are large and laminated whereas in eczema they are small and bran-like in character.



FIG. 162

Pityriasis Lichenoides

The lesions erupt as firm rounded papules which become scaly and have some resemblance to guttate psoriasis and lichen planus. Some may become pustular and leave punctate scarring. They continue to erupt over long period.



FIG. 163

Erythema Multiforme

The lesions consist of circumscribed circular infiltrated erythematous plaques each may develop bulla in the centre. Individual lesions persist for seven to ten days and the eruption lasts for about fortnight. Similar lesions are frequently present on the back of the feet, the lips, and the back of the neck.



FIG. 160

Eczema

This shows a patch of infective eczema surrounded by secondary staphylococcal pustulation.



FIG. 161

Pustular Bacterid

The lesion are vesicopustules, but the content are sterile. They tend to crust over a long period. The histopathology is distinct.



FIG. 166

Psoriatik



FIG. 166

Fig 166 A.—The lesions are fully developed and show typical psoriatic scaling.
 Fig 166 B.—The lesions have developed beneath the stratum corneum and no scaling is as yet apparent.



FIG. 167

Lapsus Erythematosus

The eruption consists of telangiectatic erythema, with slight scaling and there is an associated atrophy of the skin. The absence of an encrusted phase distinguishes it from an eczema reaction. The presence of some degree of cutaneous atrophy and the absence of large wax-like scales, from psoriasis,



FIG. 164

Psoriasis

The lesions consist of erythematous plaques, the scales are large, wax like, and heaped up on the surface in laminated layers. It may be necessary to scratch the surface to demonstrate these characters.



FIG. 165

Arthropathic Psoriasis

The nails are involved by the eruption. Waxy scaling could be demonstrated by scratching the surface of the lesions.



FIG. 170

Atrophic Lichen Planus

This is a rare form of lichen planus in which the customary aggregation of elevated flat papules is converted into an atrophic plaque. It has to be differentiated from the late stages of scleroderma and from lichen sclerosus et atrophicus. The diagnosis depends on the histopathology.



FIG. 171

Lichen Planus

The eruption consists of isolated and aggregated flat papules which are not scaly on the surface.



FIG 168

Pityriasis Rubra Pilaris

There is redness and slight scaling of the entire skin surface of the backs of the hands. The diagnostic feature is the presence of black horny plugs in the orifices of the hair follicles on the dorsum of the first phalanx.



FIG 169

Pityriasis Rubra Pilaris

There is slight hyperkeratosis with scaling. It differs from traumatic, post eczematous, and climacteric hyperkeratosis by the even formation of the excess keratin. Lesions are usually present on other areas, but the appearance is so characteristic that once it has been appreciated it is usually possible to make a diagnosis even when the palms alone are affected.



FIG. 174

Hyperkeratosis

Localized traumatic hyperkeratosis due to irregular cornification. This localized form often appears to arise spontaneously and is seen in middle and later life.



FIG. 175

Keratoderma Clavicularum

This is a mild case which somewhat resembles the lesion in toes palmaris. It is usually symmetrical, affecting both palms, the soles, and often the knees.



FIG. 172

Tinea Palmaris

The fungus has invaded the stratum corneum causing scale formation. This when removed reveals the glistening upper surface of the rete mucosum in the form of a sharply defined pink area bounded by a ribbon of scale. The condition may from time to time develop an eczema reaction.



FIG. 173

Traumatic Hyperkeratosis

This condition may follow the eczema reaction or arise spontaneously as a result of wear and tear causing the irregular formation of the stratum corneum.



FIG. 178

Nodulocystic Syphilis

The lesion consists of an aggregation of rounded nodules. It is unilateral and differs from lichen planus in which there are flat angled papules. It might suggest necrobiosis lipoidica, and could be differentiated by histopathology and serology.



FIG. 179

Scabies

Infused vesicles, persistent excoriations, and small patches of eczema, due to the presence of the scarus and to scratching.



FIG 176
Lichen Planus

Individual papules can be distinguished even in the plaque on the front of the wrist. The milky white spots and striae on the surface of the lesions are a common feature and are due to the irregular thickening of the stratum granulosum.



FIG 177
Lichen Planus

The lesion consists of an aggregation of flat non-sealy lichen planus papules. The surface is shiny due to stretching. The absence of laminated scaling and evidence of vesiculation distinguishes it from psoriasis and an eczema reaction. Typical isolated papules are usually to be found at the edges of the plaque.




FIG. 182

Paronychia

The nail bed is affected and the nails are raised from it by the acutely lesions of paronychia. The detached nail plates become friable and fragmented. In the absence of body lesions *trunc unguitum* might be suspected, and could be excluded by microscopic examination of scales and nail fragments.



FIG. 183

Ringworm of the Nails

The distal portions of the nail plates have become invaded by the fungus growth and have become fragmented.




FIG. 180

Paronychia

This shows an acute early lesion




FIG. 181

Paronychia

Here the infection has been present for some time
Note the absence of the cuticle from the affected nail.



FIG. 186

Epidermolysis Bullosa

Traumatic subepidermal bullae are present on pressure points, and the skin is atrophied in places as result of previous lesions.



FIG. 187

Epidermolysis Bullosa

T show the dental malformation frequently seen in this condition.



FIG. 184

Necrobiosis Lipoidica

The lesions resemble those of granuloma annulare, but show mild inflammation and yellowish coloration in places. They consist of flat nodules and plaques. Ulceration may occur. The histopathology is characteristic. The majority of cases occur in diabetic subjects.



FIG. 185

Granuloma Annulare

The lesions consist of pale pink or gray like nodules arranged in an annular formation. Compare necrobiosis lipoidica, from which it can be differentiated by histopathology.



FIG. 186

Epidermolysis Bullosa

Traumatic subepidermal bullae are present on pressure points, and the skin is atrophied in places as result of previous lesions



FIG. 187

Epidermolysis Bullosa

T show the dental malformation frequently seen in this condition.



FIG. 184

Necrobiosis Lipoidica

The lesions resemble those of granuloma annulare, but show mild inflammation and yellowish coloration in places. They consist of flat nodules and plaques. Ulceration may occur. The histopathology is characteristic. The majority of cases occur in diabetic subjects.



FIG. 185

Granuloma Annulare

The lesions consist of pale pink or ivory-like nodules arranged in an annular formation. Compare necrobiosis lipoidica, from which it can be differentiated by histopathology.



FIG. 190

Xanthoma

The natural folds of the palm show yellow streaks due to fat deposit.



FIG. 188

Dermatomyositis

There is streaked reticulated erythema, hypertrophy of the skin of the knuckles, and wasting of the interosseous muscles.



FIG. 189

Scleroderma

The skin is hard and shiny and movement is restricted.



FIG. 193

Milkers' Nodules

The lesions are due to infection from cows' udders, probably with cowpox virus.




FIG. 194

Fibrochondroma

This is rare benign tumour formation affecting the knuckle pads.



FIG 191
Verruca Vulgaris



FIG 192
Verruca Vulgaris



FIG. 197

Orf

This is due to virus infection contracted from sheep suffering from pustular dermatitis in the neighbourhood of the hooves. The lesion commences as an oedematous dermic infiltrate producing an inflammatory nodule. A flaccid bulla develops on the surface of the lesion.



FIG. 198

Orf

The condition commences as an oedematous infiltrate producing an inflammatory nodule. A flaccid bulla develops on its surface and eventually a crusting area is exposed.



Fig 195

Granuloma Pyogenicum

The lesion consists of granulation tissue and develops at the site of a puncture wound. Compare orf and melanoma.




Fig 196

Malignant Melanoma



FIG. 201

Diphtheritic Ulceration

Bacteriological examination of material from the surface and edge of the lesion is necessary to establish diagnosis. Systemic symptoms may develop.



FIG. 202

Blasomycosis

This rare form of fungus infection which produces chronic ulcerative granulations. Compens. tuberculosis and Bowen's disease. The causative organism (blasomycosis) can be demonstrated on culture and in histological sections of tissue.



FIG. 199

kerion

The lesion is a granuloma of short duration which develops as a result of animal ringworm infection.



FIG. 200

Leishmaniasis

Compare carcinoma and nodulocutaneous syphilide. A diagnosis is made from the macroscopic finding of Leishman Donovan bodies in serum and tissue obtained from the edge of the lesion.



FIG. 205

Basaloid Cyst

The lesion results from disorganisation of epidermal growth which represents form of intra-epidermal tumour formation. In due course the change may at some times, resolve itself into squamous cell carcinoma. The histopathology is necessary for diagnosis.



FIG. 206

Atrophy

This condition of atrophy associated with thinning of the skin (ichthyosiform), and the development of hyperkeratotic points (keratoma), usually results from prolonged exposure to sunlight and hydrocarbon derivatives.



FIG. 203

Tuberculosis Verrucosa Cutis

This form is due to external inoculation of the tubercle bacillus in an individual already sensitised to it. The reaction differs from that in lupus vulgaris by the marked hyperkeratosis which develops on the surface.



FIG. 204

Tuberculosis Verrucosa Cutis



FIG. 209

Squamous Cell Carcinoma

In this case no precarcinomatous lesions had been noted.



FIG. 207

Keratomas

The lesions in this case are due to prolonged contact with hydrocarbon derivatives and are closely allied to cutaneous horns. Malignant degeneration is prone to occur




FIG. 208

Keratomas and Carcinomas

The keratomata in this case are due to prolonged contact with shale oil, and on one hand several squamous cell carcinomata have developed from the hyperkeratotic lesions.



FIG. 212

Glomus Tumor

This is painful angioma which develops in adult life from the glomus tissue of the peripheral terminal vessels. It almost always occurs on the fingers or toes.



FIG. 213

Calcinosis Cutis

Varying sized deposits of calcium salts develop slowly in the dermis and subcutaneous tissue. Ulceration of the overlying skin and extrusion of the deposit may occur. The condition may accompany scleroderma.



FIG. 210

Myxomatous Cyst

These cysts contain a clear jelly-like liquid. They occur in the neighbourhood of the terminal phalanges, usually of the fingers. The cysts have no endothelial lining. In some cases they have been thought to communicate with the joint cavity.



FIG. 211

Myxomatous Cyst



FIG. 216

Eczema

Contact eczema of the foot. The eczematous skin has become bullous. The original vesicles can be seen as brownish points scattered over the heels. The latter is subepidermal.



FIG. 217

Impetigo

The bullous character of the impetigo is evident because of the thick stratum corneum on the sole. A similar appearance can be produced by syphilis in the newborn.



FIG. 214
Supernumerary Digit




FIG. 215
Ehlers-Danlos Syndrome

The extreme elasticity of the skin is shown, and also one of the lipomatous tumours which are commonly present in the region of the joints.



FIG. 220

Eczema

An infective (pyogenic) eczema of the toes. A similar appearance could be produced by fungus infection (*truxa pedis*), and microscopic examination of scales and vesicle roofs is necessary to exclude this.




FIG. 221

Paronychia

The scale and perionychial skin are affected.



FIG. 218

Tinea Pedis

The skin in the fourth interspace is golden, due to the growth of a fungus.



FIG. 19

Eczema

Flexural infective eczema of the toes. The appearance in Figs. 219 and 220 can be produced as a result of a fungus infection or an infection by pyogenic organisms following trauma or fissuring. Microscopic examination of scales and the roofs of vesicles is necessary procedure in diagnosis.



FIG. 224

Keratoderma Climactericum

The condition also occurs on the palms and knees.




FIG. 225

Eczema

Post-traumatic infect. eczema in dry scaly state.
The nails show onychogryphosis.



FIG. 222
Pruritus




FIG. 223

Keratoderma Blennorrhagicum (Reiter's disease)

The lesions resemble those of pruritus but are much more heaped up on the surface, and other symptoms of Reiter's disease are present.



FIG. 228



FIG. 228 a.

Subungual Exostosis

The cutaneous lesion consists of areas of granulation tissue in the region of the tip of the nail (Fig. 228). It develops as result of pressure due to an exostosis on the terminal phalanx (Fig. 228).



FIG. 226

Verruca Plantaris

The lesions are compressed into the substance of the skin and surrounded by a callosity. The surface of the verruca is rough, and small black dots are frequently to be seen scattered over it. The lesion may have to be differentiated from a localised hyperkeratosis or a corn.



FIG. 227

Taberose Sclerosis

Multiple fibromata in the region of the nails. This condition may accompany that of adenoma sebaceum.



FIG. 231

Eczema

Post-traumatic infective eczema of leg showing multiple circinate patches in close apposition. The stage of vesiculation has passed and the patches show eczema pits which are oozing droplets of plasma.



FIG. 232

Eczema

Post-traumatic infective eczema of the leg in dry scaly stage. (Compare the appearance of the reaction as shown in Fig. 242.)



FIG 229

Contact Eczema

Note the single large patches with an indefinite edge. The eruption is primarily vesicular but secondary subepidermal bullae have developed.



FIG 230

Eczema

Contact eczema of leg showing excruciation and secondary bulla formation. The margin is indefinite. In parts the epidermis forming the roofs of the vesicles has been exfoliated, revealing the bases of vesicles (eczema pits).



FIG. 235
Staphylococcal Folliculitis



FIG. 236
Pyoderma

Folliculitis and impetigo of the lower third of the leg.
This state may be the precursor of an infective eczema.



FIG. 233

Eczema

A large patch of post-traumatic infective eczema formed by the coalescence of several small patches. The reaction is in the dry scaly stage and resembles psoriasis.



FIG. 234

Eczema

A Recurrent infective eczema behind the knee which has



FIG. 235
Straphylomacul Folliculitis



F 236
Pyodermia

Folliculitis and impetigo of the lower third of the leg.
This state may be the precursor of a wider pyodermia.



FIG. 233

Eczema

A large patch of post traumatic infective eczema formed by the coalescence of several small patches. The reaction is in the dry scaly stage and resembles psoriasis.



FIG. 234

Eczema

A flexural infective eczema behind the knee which has



FIG. 239

Eczema

The right leg is affected by post-traumatic infective eczema and the exudate has formed crust which masks the reaction. The left leg shows petechial varicose hemorrhages but no eczema.



FIG. 40

Eczema

The same case as in Fig. 239. The secondary crusting has been removed from the right leg, showing vesicular



FIG 237

Eczema

A patch of infective eczema on the leg. There is also pustulation, and the eczema has probably arisen from the pustules.



FIG 238

Dermatitis Gangrenosa

This is a chronic deep-seated pyoderma usually associated with general debility



FIG. 243

Eczema

An infective eczema of the leg showing cracked appearance. This appearance of the eczema reaction is seen almost exclusively on the legs in middle age and after particularly in males.



FIG. 44

Dermatitis Asteoplytica

A simple non-eczematous dermatitis due to the application of strong detol.



FIG. 41

Eczeema

A more detailed view of the right leg shown in the previous illustration. The intense redness is in part due to vasodilatation and in part to a loss of the stratum corneum.



FIG. 42

Eczeema

A close up of the right leg in Fig. 41 to show the eczema pits.




FIG. 246

Keratoderma Climactericum

The lesions affect the fronts of the knees and the palms. They consist of hyperkeratotic plaques and are to be distinguished from the scaly lesions of psoriasis.



FIG. 47

Psoriasis

There is much scaling on the knees, but it is minimal on the hands and arms, due to frequent washing.




FIG. 245

Eczema

A patch of lichenified eczema (neurodermatitis or lichen simplex)




FIG. 246

Keratoderma Climactericum

The lesions affect the fronts of the knees and the palms. They consist of hyperkeratotic plaques and are to be distinguished from the scaly lesions of psoriasis.



FIG. 247

Psoriasis

There is much scaling on the knees, but it is minimal on the hands and arms, due to frequent washing.



FIG. 45

Eczema

A patch of lichenified eczema (neurodermatitis or lichen simplex).



FIG. 230

Porphyry



FIG. 231

Misjeck's Drift

A teluspectate and porphyric eruption which is probably synchronous in character and high is of slow development and long duration.




FIG. 248

Purpura

A mild eruption of post-influenzal purpura affecting the leg.




FIG. 249

Purpura

The eruption does not blanch on vitropression.



FIG. 254
Thlas Carpers
 (*Thlas circinata* type.)



FIG. 255
Thlas Carpers
 ("Dobolsch" type.)



FIG 252

Herpes Zoster

The eruption is profuse and consists of groups of vesicles.



FIG 253

/ Epidermolysis Bullosa

A traumatic subepidermal bulla is present and the adjacent skin shows atrophic scarring, due to previous lesions.




FIG. 257

Purpura

In this case the eruption is linear in character resembling hyperkeratotic nevus. The lesions are erythematous, and light scraping of the surface shows the wax-like scales in the illustration on the right.



FIG. 258

Pityriasis Rubra Pilaris

The eruption, which usually consists of small follicular papules, has become confluent and forms erythematous plaques which have dry rough surface. There are still islands of almost unaffected skin on which isolated papules can be seen. The eruption might suggest psoriasis or lichen planus, but waxy scales are absent and there is no infiltration such as is seen in confluent lichen planus.



FIG. 256
✓ Erythema Multiforme
Showing an annular appearance.



FIG. 261

Lichen Planus

The individual groups of papules have developed a 'starry' surface.



FIG. 262

Melanoma Cutaneous

The lower aspect thigh is a common site. In this case there is superficial resemblance to xanthomas, but the lobulated interior of the lesions can easily be expressed.



FIG. 259

Lichen Planus

In this case the process of aggregation of papules has gone further than in other examples and warty nodules have been formed (*lichen nodularis*, *lichen obtusus*). The diagnosis would depend largely on the recognition of isolated typical papules in the neighbourhood of the nodules.




FIG. 260

Lichen Planus

A chronic eruption on the leg. The papules, where single, are larger than usual but maintain their flat shiny appearance. A warty surface has developed in places on aggregations of papules.



FIG. 265

Necrobiosis Lipoidica

The eruption consists of reddish brown nodules which become aggregated to form plaques of varying shapes and sizes. In contradistinction to xanthoma there may be some scaling and ulceration may develop. Diabetes is present in the majority of cases.



FIG. 266

Necrobiosis Lipoidica

The lesions commonly affect the leg and are not infrequently linear. They may bear some resemblance to psoriasis, but they are indurated. The histopathology is diagnostic.




FIG. 263

Xanthoma Tuberosum

The elbows and knees are usual sites for this rare eruption. The yellowish nodules are smooth and non-scaly on the surface. Diabetes is almost always present.




FIG. 264

Xanthoma Tuberosum



FIG. 269

Lupus Erythematosus Profundus

This is rare form of the disease which in the absence of superficial lupus erythematosus lesions elsewhere could have to be differentiated from lupus vulgaris and nodulocystic acne.



FIG. 270

Lupus Erythematosus Profundus



FIG. 267

Lupus Vulgaris

There is some resemblance to psoriasis, but lamellar waxy scaling is absent. The plaques are slightly indurated and apple-jelly nodules can be demonstrated. Apart from the general appearance the histology would differentiate these lesions from necrobiosis lipoidica.



FIG. 268

Lupus Vulgaris

As in the previous example there is some resemblance to psoriasis, but the surface is warty and is not covered with waxy scales.



FIG. 272

Varicose Ulcer

An early lesion showing surrounding passive congestion.




FIG. 273

Varicose Ulcer

A band type of ulcer. The surrounding skin is heavily pigmented, due to passive congestion.



FIG. 71

Syphilis

Ulcerated nodulocutaneous lesions.



FIG. 276

Venous Ulcer

The lesion has healed, leaving the skin and subcutaneous tissues in a sclerosed condition.



FIG. 274

Varicose Ulcer

The surrounding skin is affected by staphylococcal folliculitis.



FIG. 75

Varicose Ulcer

The ulcer is superficial and there is a complicating infective eczema on the surrounding skin. The area is thickened, due to venous and lymph stasis.




FIG. 279

Ecdryma

The eruption is deep-seated staphylococcal infection of the hair follicles. It may arise spontaneously or as a result of contact with hydrocarbon derivatives.



FIG. 280



FIG. 281

Erythema Nodosum

The lesions are tender acute inflammatory nodules which leave temporary bruised appearances as they disappear. The lesions are always bilateral and they do not ulcerate.



FIG 277

keloid

The scar tissue is hypertrophic and in addition it enlarges at the periphery by the formation of finger-like extensions.



FIG 78

keloid




FIG. 284

Keratoma

Multiple lesions on leg. They may result from prolonged exposure to heat. A squamous cell carcinoma may develop at the base of keratoma.




FIG. 285

Carcinoma

Multiple primary lesions of the leg initiated on an old burn scar.



FIG 282

Erythema Induratum

The lesion is a slowly developing, chronic, deep-seated inflammatory nodule which usually ulcerates. Recurrence is common.



FIG 283

Erythema Induratum

The lesions may be multiple and almost always affect the lower half of the posterior aspect of the legs. They may or may not be bilateral.



FIG. 288

Mycosis Fungoides

The tumours consist of accumulations of reticulum cells in the dermis and hypoderm. Ulceration is common. Spontaneous regression takes place after variable period, to be followed by fresh crops of tumours. The diagnosis depends on the histopathology



FIG. 289

Mycosis Fungoides



FIG. 286
 Carcinoma
 Secondary cutaneous deposits.



FIG. 287
 Melanoma

The lesion is malignant in character but is only slightly pigmented. The diagnosis depends on histopathology.



FIG. 292

Hyperkeratotic Nerve

The malformation consists of a localized excess of keratin. The lesion resembles inclusionoma, but the black appearance is due to sulphur compounds in the keratin and not to melanin.



FIG. 293

Nerve Pigmentary et Pilous



FIG. 290

Dermato-fibro-sarcoma

This tumour is of low-grade malignancy. The leg is the usual site and the diagnosis depends on the histopathology.




FIG. 291

Sarcoma

(Reticulum cell type)

The diagnosis depends on the histopathology.



FIG. 295

Scabies

The lesions are scratched papules.

Figs. 295 to 302 show the eruption of scabies in varying degrees of profusion and secondary infection. The distribution of the lesions is, however uniform



✓ FIG. 296

Scabies

Some degree of secondary infection is present. Note the involvement of the elbow.




FIG 294

Glomus Tumour

Multiple lesions on upper arm. The usual sites for a glomus tumour are the fingers and toes. In this case the histopathology was necessary for diagnosis.



FIG. 299
Scabies



FIG. 300
Scabies
Detail of the impetigo on the abdomen.



✓ FIG. 297
Scabies

As a result of scratching, impetigo and folliculitis have developed.



FIG. 298
Scabies

In Figs 296 to 298 a secondary impetigo has developed.



FIG. 303

Impetigo

Ruptured and unruptured bullae are seen. The condition is primary infection, and in this form is rarely met with on this side except in children. Usually it is complication of scabies.




FIG. 301

Scabies

This illustration and that shown in Fig. 299 are of the same case. There is a severe complicating impetigo on the anterior aspect of the trunk only.



FIG. 302

Scabies

There is a profuse papula eruption on the trunk and limbs, partly due to the parasite and partly to pyogenic infection.




FIG. 306

Pityriasis Rosea

The eruption is mostly confined to the trunk and the proximal parts of the limbs.



FIG. 307

Pityriasis Rosea

The lesions consist of small discrete erythematous squamous macules, which are not infiltrated and only slightly scaly. These features differentiate it from papulosquamous syphilide and psoriasis. The roseolar syphilide is faint macular erythema, which is not scaly. An erupting pityriasis rosea may resemble rubella, but the course of the eruptions of differentiates them.




FIG. 304

Eczema

A contact eczema due to dye in clothing




FIG. 305

Psoriasis

The eruption is composed of discrete non-indurated papules which show waxy lamellar scaling.




FIG. 306

Pityriasis Rosea

The eruption is usually confined to the trunk and the proximal parts of the limbs.



FIG. 307

Pityriasis Rosea

The lesions consist of small discrete erythematous macules which are not infiltrated and only slightly scaly. These features differentiate it from a papulovesicular syphilide and psoriasis. The roseolar syphilide is faint and the erythema here is not scaly. An erupting pityriasis rosea may resemble rubella, but the course of the eruptions will differentiate them.




FIG. 308

Papulosquamous Syphilide

The papules are indurated and surmounted by a single scale. When removed this leaves a collarette of scale at the periphery of the lesion, and the exposed surface may become moist and even crusted.




FIG. 309

Papulosquamous Syphilide

The lesions are *indurated* squamous papules. The scale often becomes detached from the summit of the papule, leaving a peripheral collarette of scale.



FIG. 310

Lichen Planus

The eruption is composed of numerous slightly raised flat non-irritated and non-ecchy papules.



FIG. 311

Lichen Planus

In this case the lesions are older and somewhat pigmented.



FIG. 312

Lichen Planus

The eruption is of long-standing, and it has become deeply pigmented and slightly warty on the surface.



FIG. 313

Lichen Nidus

Compare lichen planus. The two conditions can be differentiated macroscopically. The papules are smaller than those of lichen planus, but otherwise resemble them closely. In this case a plaque has developed through the coalescence of papules. The histopathology is typical and quite distinct from lichen planus.




FIG. 314

Psoralea




FIG. 315

Psoralea



FIG. 316

Psoriasis

The silvery scales have been demonstrated by scratching
(Tâche de bougie.)




FIG. 317

Psoriasis

A large thickened plaque on which is situated a small
squamous-celled carcinoma, due to prolonged ingestion
of arsenic.



FIG. 318

Psoriasis

Either as result of confluence of papules or the central regression of plaques, annular lesions have been produced resembling tree trunks.



FIG. 319

Psoriasis

Annular form. Note that the ribbon-like lesions show typical psoriasis scaling.



FIG. 320

Tinea Corporis

The lesions bear a superficial resemblance to annular
pсориаз.



FIG. 321

Tinea Corporis

The ringed edge is composed of papules and scales, and
there are no waxy scales. Fungus elements can be
demonstrated in scrapings from the edge.



FIG. 322

Erythema Annulare Centrifugum

This is a persistent slightly infiltrated erythema which spreads centrifugally in the form of rings. Compare the gyrate erythema of dermatitis herpetiformis. The lesion is an erythema, and neither vesicles, pustules, nor scales are present.



FIG. 323

Herpes Zoster

This is a slight clue. The grouping of the vesicles, their deep-seated character, the unilateral distribution, and the attendant pain are diagnostic points.



FIG. 324
Herpes Zoster




FIG. 325
Herpes Zoster

This is a flond eruption. The unilateral character is well seen in the midline of the chest.



FIG. 326

Linear Hyperkeratotic Nerves

The lesion consists in an abnormality of keratin formation. The dark colour is due to the presence of sulphur compounds in the keratin and not to melanin.




FIG. 3.7
Urticaria




FIG. 3.28
Dermatographism



FIG. 329
Acacia Vulgaris
Comodo and superficial postular variety



FIG. 330

Oil Acne

A severe acneiform eruption in an adult, consisting of comedones, papules, and pustules, due to the action of mineral oil on the follicular orifices.




FIG. 331

Oil Acne



FIG. 332
Acne Vulgaris
Postular variety



FIG. 333
Keloid
Following acne lesions.



FIG. 334

Necrobiosis Lipoidica

There is an irregularly shaped infiltrated plaque which has undergone some atrophy in the centre (See the same condition on the legs and hands.) In the differential diagnosis of this example lupus vulgaris and syphilis would have to be considered.



FIG. 335

Lupus Vulgaris

This is an extensive uncomplicated case of many years duration. Apple-jelly nodules are easily demonstrated.



FIG. 336

Nodulocystic Syphilis



FIG. 337

Nodulocystic Acne

A few active lesions are present and there is gross scarring.



FIG. 338

Nodulocystic Acne

The lesions bear some resemblance to multiple basal cell carcinomata. Their duration, combined with serology and histopathology clarify the diagnosis.



FIG. 339
Pityriasis Versicolor



FIG. 340
Pityriasis Versicolor

The eruption due to the invasion of the stratum corneum by the *microsporum furfur*. The organism produces slight scaling, which is only evident on scratching. The affected skin has a light red or brown appearance and the condition may have to be differentiated from *scabies*. This is done by microscopic




FIG. 341

Vitiligo

The condition consists in the development of depigmented patches, surrounded by areas of hyperpigmentation. If the latter is prominent as in this case the condition at first sight might suggest pityriasis versicolor. Microscopic examination of surface scrapings will decide the point.



FIG. 342

Scleroderma

The affected area is depigmented, but the presence of sclerosis differentiates the lesion from that of vitiligo.




FIG. 343

Lichen Sclerosus et Atrophicus

Compare scleroderma. The histology differentiates this condition from lichen planus atrophicus. This condition may suggest scleroderma, but as it the lesions are not sclerosed. A scleroderma lichen has reached the atrophic stage, an atrophic lichen planus, or vitiligo are distinguished from it by histopathology



FIG. 344

Lichen Sclerosus et Atrophicus



FIG. 345

Pseudoxanthoma Elasticum

This is an atrophic change in the collagen and elastic tissue of the dermis which may develop rapidly. It is permanent and most commonly affects the upper part of the trunk. The change is diffuse and results in a slight alteration in the texture of the skin affected. It is to be differentiated from colloid pseudomilium by the fact that the latter condition occurs as small papules. It might bear a superficial resemblance to lichen sclerosus et atrophicus but is to be differentiated by its colour and histopathology



FIG. 346

Pseudoxanthoma Elasticum

The lemon-yellow colour and the alteration in texture of the affected areas are seen.



FIG. 347

Molluscum Contagiosum

The side of the trunk is common site. The contents of the lesions are easily expressed as a soft lobulated mass, and this, apart from the central umbilication, distinguishes the lesions from those of xanthoma, cysts, or lymphangioma circumscriptum.



FIG. 348

Cerebellar Nerve

The structure is fibrous. Cooper lymphangioma circumscriptum, molluscum contagiosum, and xanthoma.




FIG. 349

Melanoma

A dark lesion of this type tends to show malignant change. This example is situated on the abdomen, a not uncommon site for malignant change to occur in such lesions.



FIG. 350

Verruca Sevilis

The colour is due to hyperkeratosis. Such lesions very rarely tend towards malignant change.



FIG. 351

Neurofibromatosis

The lesions are pedunculated neurofibromata the skin overlying which tends to become hyperpigmented.



FIG. 352

Neurofibromatosis



FIG. 353
Superficial Cavernous Angioma



FIG. 354
Cavernous Angioma
The lesions are superficial, but are seldom so numerous
as in this example



FIG. 355
Sclerosing Angiosarcoma



FIG. 356
Angiosarcoma Corporis Diffusum

It is extremely rare to find angiosarcomas in such profusion. The lesions are usually limited to the scrotum and digits.




FIG. 357

Lymphangioma

Compare lymphangioma circumscriptum
and granuloma pyogenicum.



FIG. 358

Nevus Pilosus et Pigmentosus



FIG. 359

Multiple Superficial Carcinomata

The lesions are basal cell in type. They may have to be differentiated from syphilitic lesions, and possibly psoriasis. The lesion on the right buttock is fungating basal cell carcinoma, quite different in appearance from the erythematous lesions on the back.



FIG. 360

Multiple Carcinomata

The lesions are superficial carcinomatous nodules which have coalesced to form plaque. Compare psoriasis



FIG 361

Carcinoma Erythematodes

This appearance is due to the spread of cancer cells through the cutaneous lymphatic network from a tumour in the breast.



FIG. 362

Mycosis Fungoides

The eruption in this case takes the form of indurated erythematous plaques. These are produced by an infiltrate of reticulation cells in the dermis. The plaques are widely distributed over the body surface.




FIG. 363

Mycosis Fungoides

Indurated plaques are present over an extensive area. The diagnosis rests on the histopathology.



FIG. 364

Mycosis Fungoides

In this example the infiltrate in the dermis is producing circumscribed tumours of varying size many of which ulcerate

In both the plaque and tumour form the eruptive elements disappear and recur
The histopathology establishes the diagnosis.



FIG. 365

Sarcoma

This is a reticulum cell type. In contradistinction to mycosis fungoides the lesions do not tend to ulcerate or to disappear spontaneously. They have to be differentiated from those of mycosis fungoides and leukaemia.



FIG. 366

Eczema

Flexural infective type. Note the symmetry of the eruption on either side of the submammary fold and the outlying noninflammatory lesions.



FIG. 367

Eczema

Post-traumatic infective eczema following a cut.




FIG. 368

Eczema

Post-traumatic infective eczema, due to scratching in scabies.




FIG. 369

Carcinoma

Paget's disease, originating the mammary duct and breast tissue. The skin lesion consist of aggregated cutaneous nodules forming a plaque which becomes eroded on the surface. The epidermis has been invaded by duct carcinoma cells.



FIG. 370

Carcinoma

Paget's disease. The skin lesion consists of condensed intra-epidermal carcinomatous nodules the surface of which is eroded.



FIG. 371

Mycetoz Fungoides



FIG. 372

Acne Conglobata

Indolent abscesses and bridged scar formation are present. In addition the condition is usually present on the back, buttocks, scalp, and face.



FIG. 373

Hidrosadenitis

Acute recurrent abscess formation involving the sweat glands.



FIG. 374

Fox-Fordyce disease

This is a papular eruption due to the presence of an inflammatory infiltrate in the region of the apocrine sweat glands. It is persistent and extremely itchy. It may have to be differentiated from lichen planus.




FIG. 375

Flexural Infective Eczema

Note the symmetry of the lesions on either side of the inguinal fold. In *tinea cruris* the upper inner aspect of the thigh is mainly involved




FIG. 376

Tinea Cruris



FIG. 377

Hidradenitis

The condition is similar to that which is more commonly seen in the axilla. In this case, in addition to the deep sweat gland abscess involving the labium, there is a secondary folliculitis on the pubes.



FIG. 378

Eczema

Flexural selective type The various folds in the area are affected in a characteristic fashion.



FIG. 379

Nonflashes



FIG. 380

Psoriasis

Large waxy scales are present on the thigh lesions but are much less in evidence in the neighbourhood of the fold of the groin. Here the lesions might suggest flexural infective eczema, but the fold is not involved symmetrically on either side.



FIG. 381
Syphilis
Double chancre.



FIG. 382
Syphilis
Papulosquamous syphide. The lesions might have to
be differentiated from those of lichen planus.



FIG. 383

Syphilis

Papulosquamous syphilide, "mucous patch" variety



FIG. 384

Lichen Planus

The lesions might have to be differentiated from those of a papulosquamous syphilide



FIG. 315
Verruca Vulgaris



FIG. 316
Carcinoma
The tumour is squamous-celled type.



FIG 387

Angiokeratomata

The lesions consist of small capillary angiomas surmounted by a small mass of keratin. They also occur on the toes, and very occasionally as a profuse eruption on the trunk and limbs.



FIG 388

Carcinoma



FIG. 389

Erwinia

Flexural infective type. The symmetrical involvement of the fold, the sharply defined margin, and the fissure at the depth of the fold are apparent.



FIG. 390

Lymphangium Chrysanthemi

The lesions exude plasma when punctured.




FIG. 391

Verruca Vulgaris (Condyloma Accumulatum)

The lesions are the same type of common wart which affects other areas of the skin but which tend to be larger and more vascular when they develop on the navel fold and genital areas. They have to be differentiated from moist syphilitic papules.



FIG. 392

Syphilis

Papulosquamous lesions (condylomata lata). The lesions are papules which tend to be larger than usual syphilitic lesions on other areas of the body. They have to be differentiated from the common wart, which may also affect the perianal region.



FIG. 393

Acne Conglobata

The condition is chronic deep-seated pyoderma which affects the buttocks, shoulders, axillae, face, and scalp. It has some resemblance to severe acne vulgaris, but makes its appearance at later age period.



FIG. 394

Papulopustular Syphilide

This syphilitic eruption may bear a superficial resemblance to guttate psoriasis.




FIG. 395

Papulopustular Syphilide

The lesions are indurated on palpation, and this differentiates them from those of guttate psoriasis.




FIG. 396

Pityriasis

The eruption is in an early stage and consists of non-indurated squamous papules. The absence of induration differentiates them from syphilitic papules.



FIG. 397

Pityriasis Rubra Pilaris

The eruption consists of non-indurated follicular papules which give the surface a rough texture.




FIG. 394

Papulopustular Syphilide

This syphilitic eruption may bear a superficial resemblance to guttate psoriasis.




FIG. 395

Papulopustular Syphilide

The lesions are indurated on palpation, and this differentiates them from those of guttate psoriasis.



FIG. 400

Pityriasis Rubra Pilaris

The eruption has become completely confluent and resembles generalised psoriasis, but scaling is much less in evidence. In this stage pityriasis rubra pilaris may have to be differentiated not only from psoriasis but from generalised lichen planus, generalised eczema, drug eruptions, and the reticuloses.




FIG 398

Pityriasis Rubra Pilaris

The papules have become confluent, producing large areas of erythroderma, but here and there clear areas have been left which still show isolated papules.



FIG 399

Pityriasis Rubra Pilaris

The eruption is becoming confluent and presents a network of erythematous plaques.



FIG. 403

Lichen Planus

The erupt is becoming generalized and in places it is confluent. Individual papules and groups of papules are however still to be made out on the forehead. Scaling conspicuous by its absence.



FIG. 401

Psoriasis

The eruption is generalised and confluent but still shows the lamellar scaling of psoriasis.




FIG. 402

Lichen Planus

A generalised case of long standing. The lesions have become cyanotic and pigmented.



FIG. 405

Eczema

A generalized papulovesicular eczema secondary to an infective eczema of the leg



FIG. 406

Lichen Planus

The papular eruption is becoming confluent with the formation of erythematous plaques. The outlines of individual papules may still be apparent in the plaques. Vesiculation does not occur and the condition of the eruption remains constant over long periods in contrast to eczema, which may alternate between wet to dry phase



FIG. 404

Eczema

A commencing generalised eruption of papulovesicles which is secondary to a primary area of infective eczema on the leg.



FIG. 405

Eczema

A generalised papulovesicular eczema secondary to an infective eczema of the leg.



FIG. 406

Lichen Planus

The papular eruption becoming confluent with the formation of erythrodermatous plaques. The outlines of individual papules may still be apparent on the plaques. Vesiculation does not occur and the condition of the eruption remains constant over long periods in contrast to eczema, which may alternate between wet to dry phase.



FIG. 407

Eczema

The entire body surface is involved in a eczema reaction. The primary area of eczema was an infective eczema of the leg and generalisation developed suddenly presumably as a result of absorption of an allergen from the primary area.



FIG. 408

Eczema

This is generalised infective eczema, due to slow extension from areas of infective eczema affecting all the body folds. Eczema may become generalised rapidly as result of absorption from primary area of contact or infective eczema, or more slowly as a result of the slow extension and ultimate confluence of multiple patches. It may also be produced by the external administration of the heavy metals. In its dry scaly phase generalised eczema may have to be differentiated from generalised psoriasis, generalised pityriasis rubra pilaris, generalised lichen planus, and the reticuloses. The histopathology and bacteriology of the case may be necessary for diagnosis.



Fig. 409

Eczema

A generalised eczema secondary to a contact eczema on the hands.




FIG. 410

Eczema

The same case as Fig. 409



FIG. 411

Eczema

The face lesions in a case of generalised infective eczema.



FIG. 412

Eczema

The same case as is shown in F g. 411



FIG. 413

Eczema

Figs. 411 to 413 show generalised eczema secondary to an infective eczema of the ankle. The generalisation was of sudden onset.



FIG. 414

Dewler's Prurigo

The infective eczema on the breasts and the papular eruption on the chest might suggest scabies. The presence of lichenification and the history reveal the nature of the condition.



FIG. 415

Besnier's Prurigo

Note the lichenification with superimposed papulo-vesicles on the popliteal areas.



FIG. 416

Besnier's Prurigo

An acute exacerbation of the eczema reaction on lichenified skin.



FIG. 417

Eczema

A commencing Besnier prurigo. This eczema has followed an infantile eczema and in due course will become Besnier's prurigo.



FIG. 418

Dermatitis Herpetiformis

Juvenile form. A subepidermal bullous eruption. A very recent eruption might suggest varicella but the course of the disease would clarify the diagnosis. The histopathology of the bullae in the two conditions is quite distinct.



FIG. 419

✓ **Dermatitis Herpetiformis**

Juvenile form. A profuse eruption of subepidermal bullae showing a herpetiform grouping of the lesions. A similar condition can be produced by bromides or iodides.



FIG. 420

Dermatitis Herpetiformis

Juvenile form. Bunched tense subepidermal bullae. Note that the bullae are situated on uninfamed skin. This feature was formerly regarded as favouring diagnosis of pemphigus vulgaris, but it has no such significance.



FIG. 421

Dermatitis Herpetiformis

Juvenile form. The eruption is entirely composed of persistent gyrate erythematous plaques and might suggest diagnosis of erythema gyratum centrifugum. A previous history of bullae, or their subsequent appearance, would clarify the diagnosis.



FIG. 422

Dermatitis Herpetiformis

Juvenile form Gyrate persistent erythema with superimposed subepidermal bullae. The eruption might suggest linea corporis and require a microscopic examination for fungus elements.




FIG. 423

Dermatitis Herpetiformis

To show a common distribution of erythematous plaques and subepidermal bullae when the disease occurs in children. The eruption illustrated might suggest an eczema, but the essential element are erythematous plaques and bullae



FIG. 424

Dermatitis Herpetiformis

The eruption consists of a mixture of grouped subepidermal bullae and infiltrated erythematous plaques with bullae situated on them.



FIG. 425

Dermatitis Herpetiformis

The patient is aged 65 years and gravely ill. Recovery subsequently took place. The appearance of the eruption, the absence of gyrate erythema, and the gravity of the constitutional symptoms suggested a diagnosis of pemphigus vulgaris, but the bullae were subepidermal.



FIG. 426

Dermatitis Herpetiformis

In a previous attack the eruption was a purely subepidermal bullous one. In the present attack the main feature is infiltrated gyrate erythema with few bullae. The attack proved fatal.



FIG. 427

Dermatitis Herpetiformis

This is the same case as it shown in Fig. 426



FIG. 428
Dermatitis Herpetiformis



FIG. 429
Dermatitis Herpetiformis

Figs. 428 and 429 show a generalized erythematobullous eruption, but on the left shoulder a portion of the eruption shows a remarkable resemblance to herpes zoster. It will be noted, however, that the zoster-like patch crosses the middle line.



FIG. 430

Dermatitis Herpetiformis

This case shows lichenoid papules which have developed at the site of pre-existing subepidermal bullae.



FIG. 431

Dermatitis Herpetiformis

This shows the upper surface of the dermis, which has been exposed by the removal of the epidermal roof of large subepidermal bullae. The appearance suggests pemphigus vulgaris rather than dermatitis herpetiformis, but the bullae are subepidermal. The patient recovered.



FIG. 432

Dermatitis Herpetiformis

The eruption is composed uniformly of subepidermal bullae. The patient is aged 70 years and succumbed to the attack.



FIG. 433

Dermatitis Herpetiformis

The patient is aged 63 acutely ill, but recovered. The bullae are subepidermal.



FIG. 434

Pemphigus Vulgaris

A sparse eruption of intra-epidermal bullae, with a complete absence of erythema.



FIG. 435

Pemphigus Vulgaris

This shows the details of the intra-epidermal bullae. The individual lesions and groups of lesions are indistinguishable clinically from those of *dermatitis herpetiformis*.



FIG 436

Dermatitis Herpetiformis

The patient is aged 74 and later died from the disease. The eruption is uniformly bullous, but the bullae are subepidermal. The histopathology is the only criterion on which to differentiate the disease from pemphigus.



FIG 437

Pemphigus Vulgaris

The eruption is purely bullous in character, the bullae show grouping but they are intra-epidermal. The disease has responded favourably to aureomycin therapy.



FIG. 438

Pemphigus Vulgaris

This is a profuse eruption of intra-epidermal bullae. A similar clinical appearance can be presented by dermatitis herpetiformis. The patient succumbed to the disease.



FIG. 439

Pemphigus Vulgaris

The bullae are intra-epidermal, and large areas of the epidermal rete mucosae are exposed. A similar appearance can be seen in



FIG 440

Pemphigus Foliaceus

The eruption consists of flaccid subcorneal bullae. It has been present for three months. The distribution suggests lupus erythematosus, and this resemblance is accentuated when crusts are allowed to accumulate.



FIG 441

Pemphigus Foliaceus

The same case as in Fig. 440 the eruption has been present for three months. The eruption in this phase has been entitled the Seneca Usher syndrome. The distribution is that of seborrhea corporis, but the lesions are subcorneal bullae.



FIG. 442

Pemphigus foliaceus

This is the same case as in Figs. 440 and 441 and is of three months' duration. Note the involvement of the scalp by flaccid bullae.



FIG. 443

Pemphigus foliaceus

The same case as Fig. 440 nine months from its onset. The bullae are more profuse.



FIG 444

Pemphigus Foliaceus

This figure shows the detail of the eruption seen in Fig. 443. The lesion is a subcorneal bulla



FIG 445

Pemphigus Foliaceus

After eighteen months the eruption is now confluent and the complete clinical picture of pemphigus foliaceus is presented. Figs. 440 to 445 illustrated the gradual development of pemphigus foliaceus



FIG. 446

Pemphigus Follicularis

This shows generalised eruption of subcorneal bullae; the gyrate and crescentic configuration of the lesions is characteristic.



FIG 447

Pemphigus Foliaceus

This case has persisted for nine years. The eruption is past the confluent stage and is gradually becoming less in amount.



FIG. 448

Pemphigus Foliaceus

This shows the details of the subcorneal bullae of the case shown in Fig. 447



FIG. 499

Pemphigus Follaceus

This is the same case as is shown in Fig. 497. The lesions are now discrete and show crescentic outlines, which is typical.



FIG. 450

Pemphigus Follaceus

The eruption in this case has gone through the stages shown in the preceding figures. The formation of subcorneal bullae has ceased and the whole skin surface is involved. The lesions are now discrete and show crescentic outlines, which is typical.



FIG 451

Dermatomyositis

The eruption in this case bears a superficial resemblance to poikiloderma.



FIG 452

Dermatomyositis

The eruption is erythematous and telangiectatic.



FIG. 453

Dermatomyositis

Showing edema and telangiectatic erythema of the face.




FIG. 454

Dermatomyositis



FIG. 455

Dermatomyositis

The eruption consists of a telangiectatic streaked erythema.



FIG. 456

Dermatomyositis

Showing edema and erythema, there being some resemblance to acute lupus erythematosus.




FIG. 457
Pellsoderma (Atrophicum Vasculare)



FIG. 458
Pellsoderma (Atrophicum Vasculare)
Note reticulate erythema, atrophy of the skin, and patchy pigmentation. The eruption is not associated with myositis.



FIG 459

Laps Erythematosis

The eruption commenced as discrete discoid lesions on the face. It suddenly became generalised as a persistent erythema and severe constitutional symptoms developed.



FIG 460

Stevens-Johnson Syndrome

The eruption suggests a severe form of erythema multiforme, but the individual erythematous plaques and macules are atypical. It also suggests a drug eruption.



FIG. 461
Stevens-Johnson Syndrome



FIG. 462
Stevens-Johnson Syndrome
The involvement of the mucous membranes is constant feature.



FIG. 463
Herpes Zoster Varicelliformis



FIG. 464
Herpes Zoster Varicelliformis

Figs. 463 and 464 represent a severe attack of herpes zoster followed shortly after the appearance of the zoster eruption by a generalised varicelliform eruption the details of which are shown Fig. 464



A



B

FIG. 465

Acrodermatitis Chronica Atrophicans

The atrophy bears some resemblance to postuloderma but is generalized. Areas of sclerosis alternate with patches of atrophy hyperpigmentation, leucoderma, and telangiectases.



FIG. 466
Urticaria Pigmentosa



FIG. 467
Urticaria Pigmentosa

The lesions consist of persistent pigmented macules which show a transient urticarial reaction when rubbed.



FIG. 468
Darier's Disease



FIG. 469
Darier's Disease



FIG. 470
Darier's Disease



FIG. 471
Darier's Disease

When once developed this condition remains permanent with variations in degree. It consists of isolated hyperkeratotic areas which may remain discrete or become confluent and which have a typical histology. The case illustrated is a severe one and the condition may be much less extensive.




FIG. 472
Ichthyosau



FIG. 473
Ichthyosau




FIG. 474

Ichthyosis

The epidermal malformation seldom involves the flexures. Here the skin of the popliteal area is normal. This is a complete reversal of what is found in Bazier's prurigo



FIG. 475

Ichthyosis

Severe ichthyosis associated with extreme hyperkeratosis. There are all gradations of ichthyosis ranging from slight dryness and roughness of the skin to the extreme hyperkeratosis shown in this case

INDEX

A

acral tattooing, 53
 acridata, 15 16
 acrobata of aulla, 198
 acrobata, 209
 old, 35 36
 : to oil, 176
 acra, 12, 13
 acra, comedones, 175
 of face, 10, 11
 acrida, 177
 acrida, 177
 of shoulder 177
 acrida acrida and sebaceous cyst, 58
 dermatitis chronica atrophicans, 251
 acridous of hand, 100
 acrida sebaceum of face, 18, 19
 acrida in a case of, 120
 acrida acrida, 44
 acridal, 44-50
 acridal, 44-46
 acrida, 43
 acrida decaying, 43
 acrida acrida, 73
 acrida erythematosa, 46
 acrida pseudopelade of Brocq, 44
 acrida, secondary 43
 acrida, 46
 acrida capata, 47 50
 acrida acrida corporis diffusum, 189
 acrida, 206
 acrida, capillary 55, 56
 acrida, of face, 56
 of trunk, 188
 of lymphatics, 190, 208
 acrida, 189
 acrida erythema, 134, 171
 acrida, 98
 acrida, 169
 acrida, 17
 acrida, squamous carcinoma from, 168
 acrida acrida, 88
 acrida foot (Tinea pedis), 116
 acrida acrida, 223-225
 acrida acrida planus, 91
 acrida, 109

B

Bacterial, pustular 84
 Balaena, 43-50 See Alopecia
 Balaena rash (typhoid rash), 8
 Balaena cell carcinoma, cylindroma type 64, 67
 of face 64, 67 70
 multiple of face, 67
 of scalp 7
 superficial, of trunk 191
 of neck 63
 Balaena disease 190

Benign erythematoid epithelioma, 191
 Benier's prurigo, 223-225
 Blastomycosis, 107
 Blue nevus, 76
 Boeckhart impetigo (staphylococcal folliculitis), 125
 Boeck's sarcoid, 16, 17
 Botryomycosis (granuloma pyogenicum), 104
 Bowen's disease, 109
 Brocq pseudopelade, 44
 Burn scar carcinoma in, 151
 Butcher wart (tuberculous verrucosa cutis), 108

C

Calcinosus cutis, 113
 Capillary angioma, 55 56
 Carcinoma, basal cell, cylindroma type, 64, 67
 cystic, 68
 of face, 64 67 70
 multiple, 67 191
 of neck, 63
 Bowen disease, 109
 of breast, 192, 196, 197
 in burn scar 151
 erythematoid, 192
 on lupus vulgaris, 63-66
 melanotic, of face, 74 75
 of hand, 104
 non-pigmented, 152
 multiple superficial, 191
 Paget disease of the nipple, 196, 197
 in psoriasis, 168
 secondary erythematoid, 192
 nodules, 152
 squamous cell, due to arsenic, 168
 and cutaneous horn, 62
 of ear 62
 of face, 62, 64-66, 68, 69
 of hand, 110, 111
 of leg, 151
 on lupus vulgaris, 63-66
 of neck, 63
 of penis, 205, 206
 of trunk, 168
 X-ray treatment, 65-66
 Cavernous angioma of face, 56
 of trunk, 184
 Cerebelliform nevus, 185
 Character of lip, 77
 of penis, 203
 Clostrical alopecia, 44-50
 Circumscribed alopecia, 44-46
 Coal tar treatment, eczema, 85
 Cold sore (Herpes simplex), 4, 5
 Colloid pseudomycosis, 60
 Comedone acrida with sebaceous cyst, 58
 Common wart, 102

Condylomata acuminata of natal fold, 207
 of penis, 205
 lata, 208
 Contact eczema of face, 30-31
 of foot, 115
 of hand, 82
 of leg, 122
 patch test, 31
 of trunk, 162
 Contagious pustular dermatitis of sheep
 (Orf) 105
 Cracked eczema, 129
 Cutaneous horn, 61 62
 and squamous carcinoma, 62
 Cylindroma type basal cell carcinoma, 64 67
 Cyst myxomatous, 112
 sebaceous, with acneliform nevus, 58
 Cystic basal cell carcinoma, 68

D

Darier's disease, 253 254
 Deep type of lupus erythematosus, 143
 Degeneration, colloid 60
 Dermatitis. See Eczema
 autophytica, 129
 exfoliative, 218, 219
 in pityriasis rubra pilaris, 213
 in psoriasis, 214
 gangrenosa, 126
 herpetiformis, 226-234 236
 papillaris capilliti, 35 36
 seborrhoeic. See Eczema, flexural infective
 solar 32
 Dermatofibrosarcoma, 154
 Dermatomyositis, 244-246
 of hand, 100
 Dermographism, 174
 Dhobitch type of tinea, 135
 Diffuse alopecia, 43
 Digit supernumerary 114
 Diphtheritic ulceration 107
 Disease, Bazin's, 150
 Bowen's, 109
 Darier's, 253, 254
 Dühring's, 226-234 236
 Fox Fordyce, 199
 Graham-Little's, 191
 Majocchi's, 133
 Paget's, of the nipple, 196, 197
 Retter's, 118
 von Recklinghausen's, 187
 Disseminated lupus erythematosus, 248
 Drug eruption, fixed, 38
 Dühring's disease, 226-234 236

E

Ecthyma, 149
 contagiosum (Orf), 105
 Eczema, atopic, 223-225
 contact of face, 30, 31
 of foot, 115
 of hand, 82
 of leg, 122
 patch test, 31
 of trunk, 162
 cracked, 129

Eczema, flexural infective of arm, 85
 of breast, 195
 of ear 25-27
 of face, 28, 29
 of foot, 116
 of groin, 200 201
 of leg, 124
 of natal fold, 207
 of scalp, 25-28
 generalised, 218-223
 papulo-vesicular 216, 217
 infantile, 33
 transition to Besnier's prurigo, 225
 infective, of arm, 86
 of foot 117
 of leg, 126, 129
 lichenified, 37 130
 of nipple, 195
 in Besnier's prurigo, 223
 in scabies, 196
 "pits," 127 128
 post-traumatic infective, of breast
 196
 of foot, 119
 of hand, 83-84
 of leg, 123 124, 127 128
 seborrhoeic. See flexural infective
 solar 32
 tar treatment, 85
 Ehlers-Danlos syndrome, 114
 Elastosis, senile, 51
 Epidermolysis bullosa of hands, 99
 of knee, 134
 teeth in, 99
 Epiloma adenoma sebaceum in, 18 19
 nails in, 120
 Epithelioma. See Carcinoma
 benign erythematoïd, 191
 Erysipeloid 38
 Erythema annulare centrifugum, 171
 fixed, 38
 induratum, 150
 multiforme exudativum, 248 249
 of hands, 87
 of leg, 136
 nodosum, 149
 Erythematoid carcinoma, 192
 Erythrodermia in eczema, 218 219
 in pityriasis rubra pilaris, 213
 in psoriasis, 214
 Evolution of herpes zoster 6, 7
 pemphigus foliaceus, 238-240
 Exotoxins, sublingual, 121

F

Favus, 48-50
 F brochioidroma of knuckles, 103
 Fibroma of face 54
 perlingual fibrosis sclerosa, 120
 Fuliform wart 57
 Fixed drug eruption 38
 Flexural infective eczema of arm, 85
 of breast 195
 of ear 25-27
 of face, 28 29
 of foot, 116
 of groin, 200, 201
 of leg, 124

Fingert infective eczema of natal fold, 207
 of scalp, 25-28
 pruritus, 202
 Folliculitis of beard, 8
 decubitus, 45
 staphylococcal, 125
 Fox-Fordyce disease, 199
 Fungus infection of beard, 9
 blastomycosis, 107
 furva, 44-50
 of foot, 116
 of groin, 200
 of hand, 92, 106
 kerion, 9 47 48, 106
 of leg, 135
 moniliform, 202
 of nose, 97
 of neck, 9 37
 pityriasis versicolor 181
 of scalp, 47 90
 of trunk, 170

G

Generalized eczema, 216-223
 herpes zoster 134, 250
 lichen planus, 214 215
 pruritus, 214
 Gland, sebaceous, nerves of, 39-41
 sweat, nerves of, 40, 42
 Glomus tumour of arm, 156
 of hand, 113
 Graham-Little disease, 191
 Granuloma striatum, 98
 fungoides. See Mycosis fungoides
 pyogenicum, 104
 Granuloma syphilitic, 46
 Guttae pruritus, 162, 211

H

Hemangioma, capillary 35 36
 cavernous, of face, 56
 of trunk, 188
 Hair loss of 43-49
 Hair pigmented mole, 155 190
 Halo nevus, 76
 Herpes simplex, 4, 5
 lower evolution of eruption, 6, 7
 of face 6, 7
 generalized, 134
 of trunk 171, 172
 varicelliform, 290
 Hidradenitis of axilla, 198
 of perineal region, 201
 Homa, cutaneous, 61 62
 with squamous carcinoma, 6
 Homa (colloid pseudocarcinoma), 60
 Hyperkeratosis traumatic, 92, 93
 Hyperkeratosis nerves of leg, 135
 of neck, 39

I

Ichthiosis, 235, 246
 Impetigo Boeckhart (staphylococcal fol-
 liculitis), 125

Impetigo contagiosa, bullae, 115 161
 of buttocks, 161
 of face, 1 3
 of foot, 115
 Infantile eczema, 33
 transition to Besnier prurigo, 225
 Infective eczema. See Eczema

J

Jacobi's psoriasis of face, 52
 of trunk, 247
 Juvenile dermatitis herpetiformis, 226-229

K

Keloid acne, 35, 36
 following acne vulgaris, 177
 of knee, 148
 Keratoderma blennorrhagicum, 118
 characteristic of knees and hands, 131
 of palms, 93
 of soles, 119
 Kerion of face, 61
 of hand, 110
 with squamous carcinoma, 110
 of leg, 151
 leukoplakia, 80, 81
 Kerion follicularis (Darier's disease), 25
 254
 Kerion of beard, 9
 of hand, 106
 of scalp, 47 48
 Knockles, fibrochondroma of, 103

L

Leishmaniasis, 106
 Leucoderma of face and hands, 51
 of trunk, 182
 Leukoma cutis, 73
 Leukoplakia, 80, 81
 Lichen nitidus, 166
 plaques of arm, 94
 atrophicus, 91
 of buccal mucosa, 78 79
 of hand, 91
 of leg, 138 139
 of lip, 78
 of penis, 204
 pigmentation in, 166, 214
 plaque form, 94, 217
 of trunk, 166 166, 214, 19 217
 verrucous, 138, 139
 of wrist, 94
 sclerous et atrophicus, 183
 simplex of knee, 130
 of neck, 37
 Light eruption (solar eczema), 32
 Linear hyperkeratotic nevus, 173
 nevus, 39-41, 173
 pruritus, 137
 Lupus erythematosus, discoid, of face, 20,
 of hand, 89
 of lip, 79
 of scalp, 46
 disseminated, 248

-endopodite of Brocq, 44
 endothelioma elastum, 184
 keratoid, acromioid 169
 orthoplastic, 88
 carcinoma, 168
 exfoliative, 214
 flexural, 201
 of foot, 117 118
 generalized, 214
 of groin, 202
 guttae, 162, 211
 of hands, 88, 89
 of knees and hands, 131
 linear, 137
 of nails, 88, 97 117
 Tache de bougie in, 168
 of toes, 117
 of trunk, 162, 167-169, 211 214
 verrucosus (Darier's disease), 253, 254
 tumors, 152, 153
 verrucous keratoides, 133
 verrucous keratoid, 86
 verrucous, 125, 126
 verrucous granular, 104

R

"Richter's" disease, 118
 reticular cell carcinoma of face, 17
 of leg, 154
 of trunk, 194
 rhinophyma, 13
 ringworm infection of beard, 9
 of face, 42-43
 of foot, 116
 of groin, 200
 of hand, 92, 106
 of knee, 9, 47 48, 106
 of leg, 135
 of neck, 97
 of nose, 9 37
 of scalp, 47 48
 of trunk, 170
 of toes, 12, 13

S

Sarcoidosis, 14, 17
 Sarcoma, dermatofibro- 154
 of face, 72
 melanotic. See Melanocarcinoma
 reticular cell of face, 17
 of leg, 154
 of trunk, 194
 of scalp, 72
 Scarf, 157 160
 eczema of nipple in, 196
 of hand, 95
 Scalp ringworm, 47 50
 Scar carcinoma in, 155
 Scarring alopecia, 44-45
 Scleroderma of hand, 100
 of neck, 52
 of trunk, 182
 Sclerous angina, 189
 Sclerous adenoma, 18, 19
 cyst and acroform nevus, 58
 nevus, 34-41

Seborrheic eczema. See Eczema, flexural
 infective
 wart, 186
 Secondary carcinoma, 152, 192
 Senear-Usher syndrome (Pomphigus folia-
 ceus), 238-243
 Senile elastoma, 51
 wart, 186
 Shale oil keratoma and carcinoma, 110
 Shingles of face, 6, 7
 generalized, 134
 of trunk, 171 172
 varicelloform eruption, 230
 Solar eczema, 32
 Spiker nevus, 55
 Squamous cell carcinoma, due to arsenic, 168
 and cutaneous horn, 62
 of ear, 62
 of face, 62, 64-66, 68, 69
 of hand, 110, 111
 and keratoma, 110
 of leg, 151
 on lip of vulgaris, 63-66
 of neck, 63
 of penis, 205, 206
 of trunk, 168
 X-ray treatment, 65-66
 Staphylococcal folliculitis, 125
 Stevens-Johnson syndrome, 248-249
 Strawberry nevus of face, 56
 of trunk, 188
 Sublingual exostosis, 121
 Superficial cavernous angioma, 185
 Supernumerary digit, 114
 Sweat gland nevus, 40, 42
 Syphilis barbae, 8
 of face, 35 36
 Syndrome, Elderly-Darier, 114
 Senear-Usher (pomphigus foliaceus)
 238-243
 Stevens-Johnson, 248-249
 Syphilis, alopecia in, 43 46
 chancre of lip 77
 of penis, 203
 condylomata, 208
 granular, scalp, 46
 nodulocutaneous, of face, 14
 of leg, 144
 of shoulder, 179
 of trunk, 180
 of wrist, 95
 papulopustular 210
 papulosquamous, of arm, 164
 of face, 3
 of penis, 203, 204
 perianal, 208
 of trunk, 164 210
 of scalp, 43 46
 ulcerated, nodulocutaneous, 144
 Syringadenocarcinoma, 40, 42

T

Tache de bougie, 168
 Tar keratoma, 61 110
 treatment, eczema, 85
 Tattooing, accidental, 53
 Teeth in epidermolysis bullosa, 99
 Tinea barbae, 9

Lupus erythematosus profunda, 143
 pernio (sarcoid) 16
 vulgaris of arm, 142
 carcinoma on, 63-66
 of face, 22 24 179
 of leg, 142
 Lymphangioma, 190
 circumscriptum, 208
 Lymphocytoma, 60

M

Majocchi's disease, 133
 Malignant melanoma of face, 74 75
 of hand 104
 non-pigmented, 152
 tumours. See Carcinoma, sarcoma
 Marjolin's ulcer (Carcinoma in scar), 151
 Melanoma abdomen, 186
 blue naevus, 76
 malignant of face, 74 75
 of hand, 104
 non-pigmented, 152
 Menopausal keratoderma, 93 119 131
 Metastatic carcinoma, 152
 Milium, 59
 Milkers nodes, 103
 Mixed cell naevus, 54
 hyperkeratotic naevus and naevus sebaceus 39
 Mole, hairy pigmented, 155 190
 malignant. See Naevocarcinoma
 mixed cell naevus, 54
 Molluscum contagiosum of thigh, 139
 of trunk, 185
 fibrosum (neurofibromatosis) 187
 Monilliasis, 202
 Multiform erythema of hands, 87
 of leg, 136
 Multiple basal cell carcinoma, 67 191
 superficial carcinoma, 191
 Mycosis fungoides of breast, 197
 of head, 73 74
 of legs, 153
 of trunk, 193 194
 Myxomatous cyst 112

N

Naevocarcinoma of face, 74 75
 of hand, 104
 non-pigmented, 152
 Naevus, acneliform and sebaceous cyst 58
 araneus, 55
 blue, 76
 cerebelliform, 185
 halo, 76
 hyperkeratotic, of leg, 155
 of neck, 39
 of trunk 173
 linear 39-42
 wart 39, 173
 mixed cell 54
 pigmentosus et pilosus, 155 190
 sebaceous, 39-41
 syringadenomatous papilliferus, 40-42
 vascular 55 56
 wart 39 155 173

Neurobiosis lipoides of hand, 98
 of leg, 141
 of trunk, 178
 "Nettle rash," 174
 Neurodermatitis of knee, 130
 of neck, 37
 Neurofibromatosis, 187
 Nipple, Paget's disease of, 196, 1
 Nodes, milkers 103
 Nodose erythema, 149
 Nodulocutaneous syphilide of face
 of leg, 144
 of shoulder, 179
 of trunk 180
 of wrist, 95

O

Oil acne, 176
 Onychogryphosis, 119
 Orf of arm, 105
 of hand 105

P

Paget's disease of the nipple, 196
 Palmar xanthoma, 101
 Papulonecrotic tuberculide, 15, 16
 Papulopustular syphilide of trunk
 Papulosquamous syphilide of arm
 condylomata, 208
 of face 3
 of penis, 203 204
 of trunk, 164 210
 Paronychia, 96
 Patch test, 31
 Pemphigus foliaceus, 238-243
 evolution of 238-240
 neonatorum (impetigo) 115
 vulgaris, 235-237
 Perionychia, 96
 Perianal fibromata in tubercosis 120
 Pigmented mole, 155 190
 Pityriasis lichenoides, 87
 rosea, 163
 rubra pilaris of face, 34 212
 generalised, 213
 of hand, 90
 of leg, 137
 of trunk 211 213
 verrucular 181
 Plantar wart 120
 Poikiloderma atrophicum vasculare 52
 of trunk 247
 Poikilodermatomyositis, 100, 244-
 "Port wine stain," 55 56
 Post-febrile alopecia, 43
 Post-mortem wart (tuberculosis) 108
 Post-traumatic infective eczema of
 foot, 119
 of hand, 83 84
 of leg, 123 124 127 128
 Powder mark, 53
 Prurigo, Besnier's, 223-225

endopiede of Brocq, 44
 elastica elastica, 184
 eosinophilic, 169
 erythematous, 214
 carcinoma, 168
 exfoliative, 214
 flexural, 202
 of foot, 117, 118
 generalized, 214
 of groin, 202
 guttate, 16, 211
 of hands, 88, 89
 of knees and hands, 131
 linear, 137
 of nails, 83, 97, 117
 Tache de bougie in, 168
 of toes, 117
 of trunk, 162, 167-169, 211, 214
 keratogenous (Darier disease), 253, 254
 urticaria, 132, 133
 verrucous keratogenous, 133
 verrucous keratoid, 86
 verrucous, 125, 126
 verrucous granular, 104

R

Racemose disease, 118
 reticular cell carcinoma of face, 17
 of leg, 154
 of trunk, 194
 rhinophyma, 13
 ringworm infection of beard, 9
 virus, 48-50
 of foot, 116
 of groin, 200
 of head, 92, 106
 kerion, 9, 47, 48, 106
 of leg, 135
 of nails, 97
 of neck, 9, 37
 of scalp, 47-50
 of trunk, 170
 Rosacea, 12, 13

S

Sarcomatous, 16, 17
 Sarcoma, dermatofibro- 154
 of face, 72
 melanotic. See Neurocarcinoma
 melanotic cell of face, 17
 of leg, 154
 of trunk, 194
 of scalp, 72
 Sebacea, 157, 160
 eczema of nipple in, 196
 of hand, 95
 Scalp ringworm, 47-50
 Scar carcinoma to, 151
 Scarring alopecia, 44-50
 Scleroderma of hand, 100
 of neck, 52
 of trunk, 182
 Sclerotic angoma, 189
 Sebaceous adenoma, 18, 19
 cyst and acroform nevus, 58
 virus, 39-41

Seborrheic eczema. See Eczema, flexural
 infective
 wart, 186
 Secondary carcinoma, 152, 197
 Senear-Usher syndrome (Pemphigus foliaceus), 238-243
 Senile elastosis, 51
 wart, 186
 Shale oil keratoma and carcinoma, 110
 Shingles of face, 6, 7
 generalized, 134
 of trunk, 171, 177
 varicelliform eruption, 290
 Solar eczema, 32
 Spider nevus, 55
 Squamous cell carcinoma, due to arsenic, 168
 and cutaneous horn, 6
 of ear, 62
 of face, 62, 64-66, 68, 69
 of hand, 110, 111
 and keratoma, 110
 of leg, 151
 on hepato vulgaris, 63-66
 of neck, 63
 of penis, 205, 206
 of trunk, 168
 X-ray treatment, 65-66
 Staphylococcal folliculitis, 125
 Stevens-Johnson syndrome, 248-249
 Strawberry nevus of face, 56
 of trunk, 188
 Subungual exostosis, 121
 Superficial cavernous angioma, 188
 Supernumerary digit, 114
 Sweat gland nevus, 40, 42
 Syphilis barbae, 8
 nucha, 35, 36
 Syndrome, Ehlers-Danlos, 114
 Senear-Usher (pemphigus foliaceus),
 238-243
 Stevens-Johnson, 248-249
 Syphilis, alopecia in, 43, 46
 chancre of lip 77
 of penis, 203
 condylomata, 208
 gummatous, scalp, 46
 nodulocystic, of face, 14
 of leg, 144
 of shoulder, 179
 of trunk, 180
 of wrist, 95
 papulopustular, 210
 papulosquamous, of arm, 164
 of face, 3
 of penis, 203, 204
 peri-anal, 208
 of trunk, 164, 210
 of scalp, 43, 46
 ulcerated, nodulocystic, 144
 Syringocystomatous nevus, 40, 42

T

Tache de bougie, 168
 Tar keratoma, 61, 110
 treatment, eczema, 85
 Tattooing, accidental, 53
 Teeth in epidermolysis bullosa, 99
 Tinea barbae, 9

Lupus erythematosus profunda 143
 pernio (sarcoid), 16
 vulgaris of arm, 142
 carcinoma on, 63-66
 of face, 22 24 179
 of leg, 142
 Lymphangioma, 190
 circumscriptum, 208
 Lymphocytoma, 60

M

Majocchi's disease, 133
 Malignant melanoma of face, 74 75
 of hand, 104
 non-pigmented 152
 tumours. See Carcinoma, sarcoma
 Marjolin's ulcer (Carcinoma in scar), 151
 Melanoma, abdomen, 186
 blue nevus, 76
 malignant of face 74 75
 of hand, 104
 non-pigmented, 152
 Menopausal keratoderma, 93 119 131
 Metastatic carcinoma, 152
 Milium, 59
 Milkers' nodes, 103
 Mixed cell nevus, 54
 hyperkeratotic nevus and nevus sebaceus
 39
 Mole, hairy pigmented, 155 190
 malignant. See Naevocarcinoma
 mixed cell nevus, 54
 Molluscum contagiosum of thigh 139
 of trunk, 185
 fibrosus (neurofibromatosis), 187
 Moniliasis, 202
 Multiform erythema of hands, 87
 of leg, 136
 Multiple basal cell carcinoma, 67 191
 superficial carcinoma 191
 Mycosis fungoides of breast, 197
 of head, 73 74
 of legs, 153
 of trunk 193 194
 Myxomatous cyst, 112

N

Naevocarcinoma of face, 74 75
 of hand, 104
 non-pigmented, 152
 Nevus, acneliform and sebaceous cyst, 58
 araneus, 53
 blue, 76
 cerebelliform, 185
 halo, 76
 hyperkeratotic, of leg, 155
 of neck, 39
 of trunk 173
 linear 39-42
 wart 39 173
 mixed cell, 54
 pigmentosus et pilosus, 155 190
 sebaceous, 39-41
 syngangiomatous papilliferus, 40-42
 vascular 55 56
 wart 39 155 173

Necrobiosis lipoidica of hand, 98
 of leg, 141
 of trunk 178
 "Nettle rash," 174
 Neurodermatitis of knee, 130
 of neck, 37
 Neurofibromatosis, 187
 Nipple, Paget's disease of 196, 19
 Nodes, milkers' 103
 Nodose erythema, 149
 Nodulocutaneous syphilide of face
 of leg, 144
 of shoulder 179
 of trunk, 180
 of wrist, 95

O

Oil acne, 176
 Onychogryphosis, 119
 Orr of arm, 105
 of hand, 105

P

Paget's disease of the nipple, 196
 Palmar xanthoma, 101
 Papulonecrotic tuberculide, 15 16
 Papulopustular syphilide of trunk
 Papulosquamous syphilide of arm
 condylomata, 208
 of face 3
 of penis, 203 204
 of trunk 164 210
 Paronychia 96
 Patch test, 31
 Pemphigus foliaceus, 238-243
 evolution of 238-240
 neonatorum (impetigo), 115
 vulgaris, 235-237
 Perionychia, 96
 Periungual fibromata in tuberose
 120
 Pigmented mole, 155 190
 Pityriasis lichenoides, 87
 rosea 163
 rubra pilaris of face 34 212
 generalised 213
 of hand, 90
 of leg, 137
 of trunk 211 213
 versicolor 181
 Planter wart, 120
 Poikiloderma atrophicum vasculare
 52
 of trunk 247
 Poikilodermatomyositis, 100, 244-2
 Port wine stain, 55 56
 Post febrile alopecia, 43
 Post mortem wart (i. berculosa) er
 108
 Post traumatic infective eczema of
 195 196
 of foot, 119
 of hand, 83 84
 of leg, 123 124 177 128
 Powder mark 53
 Prurigo Besnier's, 223-225

INDEX

Tinea capitis, 47 50
 circinata, 135
 corporis, 37 135 170
 cruris, 200
 favosa, 48-50
 kerion of beard, 9
 of hand, 106
 of scalp, 47-48
 of nails, 97
 of neck, 9
 palmaris, 92
 pedis, 116
 versicolor 181

Traumatic hyperkeratosis, 92, 93
 Tuberculide, papulonecrotic, 15 16
 Tuberculosis, lupus vulgaris, carcinoma in,
 63-66
 of face, 22 24 179
 of leg, 142
 verrucosa cutis, 108
 Tuberosc sclerosis, adenoma sebaceum in,
 18 19
 perungual fibromata in, 120
 Tumour glomus, of arm, 156
 of hand, 113
 turban, 67 72
 Turban tumour 67 72

U

Ulcer diphtheritic, 107
 Marjolin's (carcinoma in scar) 151
 syphilitic, 144
 varicose, 145-147
 Urticaria, 174
 pigmentosa, 252

V

Varicelliform eruption in zoster 250
 Varicose ulcer 145-147
 Verruca filiformis, 57
 plantaris, 120
 senilis, 186
 vulgaris of hand, 102
 of natal fold, 207
 of penis, 205

Verrucous naevus, 39 155 173
 Vitiligo of face and hands, 51
 of trunk, 182
 von Recklinghausen's disease 187

W

Wart, common, 102
 filiform, 57
 of natal fold, 207
 of penis, 205
 plantar 120
 post-mortem (tuberculosis cutis), 108
 senile, 186
 Warty naevus of leg, 155
 linear 39 173
 of neck, 39
 of trunk 173

X

Xanthelasma, 57
 Xanthoma of palm, 101
 tuberosum of elbows and knees, 140
 of neck, 36
 papules of 140
 Xeroderma, 255, 256
 X ray photograph of subungual exostosis,
 121
 squamous carcinoma following, 63 64
 treatment of squamous carcinoma with
 65 66

Y

Yeast infection (moniliasis) 202

Z

Zoster of face, 6, 7
 generalised, 134
 of trunk, 171 172
 varicellosis, 250

